

## ASSESSMENT OF Pb, Cd, Cu AND Zn AVAILABILITY FOR PLANTS IN BAIA MARE MINING REGION

LEVEI ERIKA-ANDREA<sup>1</sup>, MICLEAN MIRELA<sup>1</sup>, ŞENILĂ MARIN<sup>1</sup>, CADAR OANA<sup>1</sup>, ROMAN CECILIA<sup>1</sup>, MICLE VALER<sup>2</sup>

**Abstract.** In order to evaluate the mobility of heavy metals in soil from Baia Mare mining region, the total, water and DTPA extractable metal contents were determined. The results showed that despite the high total metals contents and the high percentages of plant available metals only a low percent was water soluble, indicating a potential accumulation of metals in trophic chain and a potential risk for public health. Among the investigated metals, the plant available Pb and Cd species are the most severe contaminants. Significant correlations between total and DTPA extractable metals were found for Cu ( $r=0.510$ ) and Pb (0.418), and also an affinity between total and water extractable metals were identified for Cu (0.366), Pb (0.502) and Zn (0.597).

**Key words:** Heavy metals availability, DTPA extraction, soil, mining

### Introduction

Heavy metals represent a potential hazard to humans and environment. In industrial areas the heavy metal contents from anthropogenic sources are several times higher than those from natural ones [NRIAGU & PACYNA, 1988]. In addition, areas far from industrial centres also show increasing heavy metal concentrations due to long-range atmospheric transport [CEMEK & KIZILKAYA, 2006]. Mining and ore processing industry is one of the major sources of metals releasing into the environment [CASSELLA & al., 2007; BOUGHRIET & al., 2007; VANDERLINDEN & al., 2006; VANEK & al., 2005; CONESA & al., 2007; MACKLIN & al., 2003, LEVEI & al., 2009, MICLEAN & al., 2009].

Although the total metal concentration is commonly used in soil quality standards, it provides no information regarding the metals chemical nature or mobility [SILVEIRA & al., 2006; WALTER & al., 2006]. To assess the metals availability from soil to plant a great variety of single or sequential extraction schemes have been developed [FUENTES & al., 2004; PEREZ-SANTANA & al., 2007; ABOLLINO & al., 2006]. The DTPA (diethylenetriaminepentaacetic acid) extraction method was initially designed to predict micronutrient deficiencies in neutral to calcareous soils [LINDSAY & NORWELL, 1978], but it has been also employed for the estimation of metal availability for plants [MAIZ & al., 2000]. Water soluble metals represent the most ecotoxicologically relevant fraction in the environment, due to their high contamination potential of the food chain, surface water and groundwater [MEERS & al., 2006].

The objective of this study was to investigate the availability of Cu, Pb, Zn and Cd from soil to plants in Baia Mare mining region using selective extraction in water and DTPA.

<sup>1</sup> INCDO-INOE Research Institute for Analytical Instrumentation, Cluj-Napoca, 67 Donath, 400293, Romania, icia@icia.ro

<sup>2</sup> Technical University of Cluj-Napoca, 15 Constantin Daicoviciu, Romania

## Material and methods

### Site description and sampling

In the Baia Mare area, around an industrial complex involved in mining, metallurgical and chemical activities, the environment and particularly the soils are polluted due to the acid rains and heavy metal emissions [LACATUSU & al., 2001]. The region became of international concern after the cyanide spill accident in January 2000 that affected the ecosystem of Tisa and Danube rivers. Despite the fact that the large industrial plants have reduced their activities and some of them were closed, the area is still highly polluted [LEVEI & al., 2009].

A number of 50 surface soil samples were collected in the summer of 2009 from the Baia Mare town, using a stainless steel shovel. Samples were air dried to constant weight, sieved through the 2 mm sieve. The fraction < 2 mm was stored in polyethylene bags until the determination of total, water and DTPA-extractable metal contents.

### Chemical analysis

The total metal concentrations in soils were determined after aqua regia digestion. An amount of 1 g soil sample was weighted, introduced into the reaction flask and heated under reflux conditions for 2 h with 21 ml of 12 M HCl and 7 ml of 15.8 M HNO<sub>3</sub>. The solution was filtered and diluted to 100 ml with 0.5 M HNO<sub>3</sub>. Total heavy metal concentrations were determined by inductively coupled plasma atomic emission spectrometry (ICP-AES) using the scanning spectrometer SPECTROFLAME (Spectro Analytical Instruments, Kleve, Germany).

The plant available metal contents of soils were determined by extraction in water and DTPA. The DTPA extractable metal content was determined using a mixture of 0.005 mol L<sup>-1</sup> DTPA, 0.01 mol L<sup>-1</sup> CaCl<sub>2</sub>, 0.1 mol L<sup>-1</sup> triethanolamine (TEA) with pH adjusted to 7.3 with 1 mol L<sup>-1</sup> HCl solution. An amount of 10 g of soil sample was weighted into a 125 mL flask, and shaken for 2 h at room temperature using a magnetic shaker with 20 mL of DTPA extracting solution. The extracts were filtered and diluted to 100 ml with ultrapure water. The water extractable metal content was determined in a 1:10 soil/water mixture. An amount of 5 g of soil sample was shaken with 50 ml water for 2 h at room temperature. The solution was filtered and diluted to 50 ml with ultrapure water. Water and DTPA-extractable metal contents were determined by inductively coupled plasma mass spectrometry (ICP-MS) using the ELAN DRC II-Perkin Elmer, USA).

Soil pH was measured in a suspension of 1:5 soil/water ratio. The suspension was allowed to stand 1 h prior to pH measurement using a Jenway ion-meter.

## Results and discussion

The statistic parameters of total, water and DTPA extractable Cu, Pb, Zn and Cd contents are presented in Tab. 1. Due to the fact that soils were sampled from the Baia Mare town the results were compared with the guidelines values for sensitive soil according to Romanian legislation [Ministerial Decree no. 756/1997].

The total contents of all metals were high, exceeding the alert values for sensitive soils (residential and agricultural use) for the most samples. In case of Pb and Zn the 1<sup>st</sup> quartile is higher and in case of Cu and Cd is around the corresponding alert level for

sensitive soil. For all samples the 3<sup>rd</sup> quartile was much higher than the action trigger value for all metals. The average total metal content exceeded the action trigger values for sensitive use according to Romanian legislation, 1.5 times in case of Cu and Cd, 3 times in case of Zn and 18 times in case of Pb.

**Tab. 1.** Descriptive statistics of total, water and DTPA extractable Cu, Pb, Zn and Cd in soil

	Cu			Pb		
	Total	DTPA	Water	Total	DTPA	Water
	(mg kg <sup>-1</sup> )					
<b>Min</b>	38.1	6.5	1.4	87.8	0.1	0.7
<b>Max</b>	1770	187	147	23300	753	16.2
<b>Average</b>	314	58	12.5	1790	153	3.3
<b>Median</b>	168	45.3	6.9	496	87	1.6
<b>Skewness</b>	2.68	1.63	5.39	4.66	2.99	2.32
<b>Kurtosis</b>	9.02	2.32	30.2	23.3	10.4	5.23
<b>1 st quartile</b>	109	25.6	5.5	404	70	1.2
<b>3 rd quartile</b>	367	70.0	11.0	1420	156	3.5
<b>Alert value</b>	100	-	-	50	-	-
<b>Action trigger value</b>	200	-	-	100	-	-
	Zn			Cd		
	Total	DTPA	Water	Total	DTPA	Water
	(mg kg <sup>-1</sup> )					
<b>Min</b>	109	16.3	0.66	1.9	0.28	0.01
<b>Max</b>	11500	351	355	29.9	9.4	2.2
<b>Average</b>	1828	96.9	26.7	7.9	2.3	0.56
<b>Median</b>	737	75.4	6.12	3.5	1.7	0.14
<b>Skewness</b>	2.43	1.94	4.26	1.59	2.30	1.68
<b>Kurtosis</b>	5.67	3.43	19.2	2.00	6.10	2.50
<b>1 st quartile</b>	436	39.6	1.8	3.1	1.2	0.1
<b>3 rd quartile</b>	1705	118	10.9	11.4	2.8	0.9
<b>Alert value</b>	300	-	-	3	-	-
<b>Action trigger value</b>	600	-	-	5	-	-

Total = concentration of metal extracted in aqua regia

DTPA = concentration of metal extracted in DTPA

Water = concentration of metal extracted in water

The total metal contents in soil were, in all cases, higher than those from the vicinity of Pb and Ag processing smelter in the Příbram region in the Czech Republic [VANEK & al., 2005]. The median values of total Pb, Zn and Cd were similar, whilst that of Cu was lower than those from Deer Lodge valley smelter area in Montana, USA [BURT & al., 2003]. The water soluble metal contents and the average percent of DTPA-extractable Cu were higher in our study, while the average total Cu concentration was similar to that found in heavily polluted soils from the vicinity of a deactivated mining site in the Amazon region of Brazil [CASSELLA & al., 2007]. The percentages of water extractable Pb and Zn in our soil samples were similar to those found in soils from a Spanish mining region, but the average percentage of the found DTPA extractable metals was much higher [CONESA & al., 2007].

The content of DTPA-extractable Pb, Cu and Cd from soil exceeded the alert level in 83%, 9% and 17% of samples, respectively. The concentration limits of 20 mg kg<sup>-1</sup> and

70 mg kg<sup>-1</sup> DTPA-extractable Pb and Zn respectively, advocated to avoid human risk [WINTER SYDNOR & REDENTE, 2002] were exceeded in 94% and 54% of samples in case of Pb and Zn respectively. In case of Pb the 1<sup>st</sup> quartile of the DTPA-extractable content exceeded the alert value and the 3<sup>rd</sup> quartile the action trigger value for sensitive soils. A relatively high fraction of the total metal content was extracted in DTPA. Thus, an average of 22% of the total Pb content was DTPA-extractable and only 0.5% was water-extractable. The corresponding percentages for the other metals were: 11% and 2.3% for Zn, 24% and 4.4% for Cu and 43% and 4.1% for Cd, respectively.

The water soluble content of Cu, Pb and Zn was small compared to the total metal content and that of Cd was below the detection limit (1.4 mg kg<sup>-1</sup>) of the method in the majority of samples. The average percentages of DTPA and water soluble metals are presented in Fig. 1.




Fig. 1. The average percentage of DTPA and water extractable Cu, Pb, Zn and Cd

Significant correlations between total and DTPA extractable metals were found only for Cu and Pb suggesting that the percent of available metal depends on the nature and mobility of metals species exhausted by polluters. Significant correlations were found between total and water extractable metal content for Cu, Pb and Zn (Tab. 2).

Tab. 2. Correlation coefficients between total-DTPA extractable contents and total-water extractable contents of metals

	Total Cu	Total Pb	Total Zn	Total Cd
DTPA-extr.	0,510**	0,418*	0,258 <sup>ns</sup>	0,260 <sup>ns</sup>
Water-extr.	0.366*	0.502**	0.597**	0.131 <sup>ns</sup>
ns-not significant; * correlation significant at p<0.05; ** correlation significant at p<0.01				

The pH values in soil ranged between 3.0-8.23, with average value 6.74, but no statistically significant relationship between pH and available metal content was found. Strong negative correlations were found between water-extractable Cu ( $r = -0.393$ ), Zn ( $r =$

-0.532), Cd ( $r = -0.744$ ) and pH. The lack of pH influence on DTPA extractable metal contents is probably due to high buffering capacity of DTPA solution.

### Conclusions

The results showed that metal concentrations of polluted soils varied widely, in most cases exceeding the corresponding alert levels, indicating a severe situation, needing urgent measurements of pollution stopping and applying soil decontamination solutions, especially they cannot be degraded or destroyed. The high percentages of DTPA - extractable metals indicate an anthropogenic pollution, and a potential metal accumulation in vegetables, posing a potential risk for public health.

### Acknowledgements

The financial support provided by the Romanian Ministry of Education and Research, PNCDI II Project RESOLMET no. 32161/2008 is greatly appreciated.

### References

1. ABOLLINO O., GIACOMINO A., MALANDRINO M., MENTASTI E., ACETO M. & BARBERIS R. 2006. Assessment of metal availability in a contaminated soil by sequential extraction, *Water Air Soil Pollut.*, **137**: 315-338.
2. BOUGHRIET A., PROIX N., BILLON G., RE COURT P. & OUDDANE B. 2007. Environmental impacts of heavy metal discharges from a smelter in Deule-canal sediments (Northern France) concentration levels and chemical fractionation, *Water Air Soil Pollut.*, **180**: 83-95.
3. BURT R. WILSON M. A., KECK T. J., DOUGHERTY B. D., STROM D. E. & LINDAHL J. A. 2003. Trace element speciation in selected smelter-contaminated soils in Anaconda and Deer Lodge Valley, Montana, USA, *Adv. Environ. Res.*, **8**: 51-67.
4. CASSELLA R. J., WAGENER A. L.R., SANTELLI R. E., WAGENER, K. & TAVARES L. Y. 2007. Distribution of copper in the vicinity of a deactivated mining site at Carajas in the Amazon region of Brazil, *J. Hazardous Mat.*, **142**: 543-549.
5. CEMEK B. & KIZILKAYA R. 2006. Spatial variability and monitoring of Pb contamination of farming soils affected by industry, *Environ. Monit. Assess.*, **117**: 357-375.
6. CONESA H.M., FAZ A. & ARNALDOS R. 2007. Initial studies for the phytostabilization of a mine tailing from the Cartagena-La Union Mining District (SE Spain), *Chemosphere*, **66**: 38-44.
7. Decree of Ministry of waters, forest and environment protection for the approval of Regulation regarding the environment pollution assessment, no. 756/1997.
8. FUENTES A., LLORENS M., SAEZ J., SOLER A., AGUILAR M. I., ORTUNO J.F. & MESEGUER V.F. 2004. Simple and sequential extractions of heavy metals from different sewage sludges, *Chemosphere*, **54**: 1039-1047.
9. LACATUSU R., DUMITRU M., RISNOVEANU I., CIOBANU C., LUNGU M., CARSTEA S., KOVACSOVICS B. & BACIU C. 2001. Soil pollution by acid rains and heavy metals in Zlatna region, Romania, 817-821, In D. E. Stott, R. H. Mohtar and G. C. Steinhardt (eds.). Sustaining the Global Farm. Selected papers from the 10<sup>th</sup> International Soil Conservation Organisation Meeting, May 24-29, 1999 at Purdue University and the USA-ARS National Soil Erosion Research Laboratory.
10. LEVEI E., FRENTIU T., PONTA M., SENILA M., MICLEAN M., ROMAN C. & CORDOS E. 2009. Characterisation of soil quality and mobility of Cd, Cu, Pb and Zn in the Baia Mare area Northwest Romania following the historical pollution, *Internat. J. Environ. Anal. Chem.*, **89** (8): 635-649.
11. LINDSAY W. L. & NORWELL W. A. 1978. Development of a DTPA soil test for zinc, iron, manganese and copper, *Soil Sci. Soc. Am. J.*, **42**: 421-428.
12. MACKLIN M.G., BREWER P.A., BALTEANU D., COULTHARD T. J., DRIGA B., HOWARD A.J. & ZAHARIA S. 2003. The long term fate and environmental significance of contaminant metals released by the January and March 200 mining tailings dam failure in Maramures County, upper Tisa basin, Romania, *Appl. Geochem.*, **18**: 241-257.

#### ASSESSMENT OF Pb, Cd, Cu AND Zn AVAILABILITY FOR PLANTS IN BAIJA MARE ...

---

13. MAI, I., ARAMBARRI I., GARCIA R. & MOLLAN E. 2000. Evaluation of heavy metal availability in polluted soils by two sequential extraction procedures using factor analysis, *Environ. Pollut.*, **110**: 3-9.
14. MEERS E., DU LAING G., UNAMUNO V. G., LESAGE E., TACK F. M. G. & VERLOO M. G. 2006. Water Extractability of Trace Metals from Soils: Some Pitfalls, *Water Air Soil Pollut.*, **176**: 21-35.
15. MICLEAN M., LEVEI E.A., SENILA M., ROMAN C. & CORDOS E. 2009. Assessment of Cu, Pb, Zn and Cd availability to vegetable species grown in the vicinity of tailing deposits from Baia Mare area, *Rev. Chim.*, **60** (1): -5.
16. NRIAGU J. O. & PACYNA J. M. 1988. Quantitative assessment of worldwide contamination of air, water and soils by trace metals, *Nature*, **333**: 134 – 139.
17. PEREZ-SANTANA S., POMARES ALFONSO M., VILLANUEVA TAGLE M., PENA ICART M., BRUNORI C. & MORABITO R. 2007. Total and partial digestion of sediments for the evaluation of trace element environmental pollution, *Chemosphere*, **66**: 1545-1553.
18. SILVEIRA M. L., ALLEONI L. R. F., O'CONNOR G. A. & CHANG A. C. 2006. Heavy metal sequential extraction methods-a modification for tropical soils, *Chemosphere*, **64**: 1929-1938.
19. VANDERLINDEN K., ORDONEZ R., POLO M. J. & GIRALDEZ J. V. 2006. Mapping residual Pyrite after a mine spill using non co-located spatiotemporal observations, *J. Environ. Qual.*, **35**: 21-36.
20. VANEK A, BORUVKA L., MIHALJEVIC M. & KOMAREK M. 2005. Mobility of lead, zinc and cadmium in alluvial soils heavily polluted by smelting industry, *Plant Soil Environ*, **51**: 316-321.
21. WALTER I., MARTIMEZ F. & CALA V. 2006. Heavy metal speciation and phytotoxic effects of three representative sewage sludges for agricultural uses, *Environ. Pollut.*, **139**: 507-514.
22. WINTER SYDNOR M. E. & REDENTE E. F. 2002. Reclamation of high-elevation, acidic mine waste with organic amendments and topsoil, *J. Environ. Qual.* **31**: 1528–1537.