

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE / NUMBER AND ROOT ARCHITECTURE IN ROOT TIPS OF *ARABIDOPSIS THALIANA* SEEDLINGS. CONNECTIONS TO THE ROOT HAIR PATTERNING AND ABUNDANCE

Iva MCCARTHY-SUÁREZ

Hormone Signalling and Plant Plasticity Group. Instituto de Biología Celular y Molecular de Plantas (IBMCP-UPV-CSIC). Ingeniero Fausto Elio s/n (46022), Valencia – Spain.
E-mail: ivmcsua@upvnet.upv.es

Abstract: A previous study showed that excessive GAs/DELLAs altered the arrangement, shape and frequency of hairs in root tips of *A. thaliana* seedlings by acting from the root sub-epidermal tissues. The present study showed that excessive GAs/DELLAs also modified root tip cell size/number in *A. thaliana* seedlings. While excessive DELLAs shortened and widened the root epidermal, cortical, endodermal and pericycle cells, excessive GAs, excepting the epidermal cells, generally narrowed them. However, no root cell size changes occurred when *gai-1* was over-expressed at the root epidermis. Also, excessive DELLAs induced extra cells at the root epidermis, cortex, endodermis and pericycle and lateral root outgrowth, whereas excessive GAs induced extra cells at the root cortex and pericycle. Thus, excessive GAs/DELLAs might modify root tip cell size/number by acting from the sub-epidermal tissues of the root. This, in turn, might impact on the root hair patterning & abundance and the root architecture.

Keywords: DELLAs, Gibberellins, lateral roots, root architecture, root cell number, root cell size.

Introduction

A previous study showed that supra-physiological levels of GAs/DELLAs altered the patterning, the morphology and the abundance of root hairs in *A. thaliana* seedlings, and that they did it by possibly acting from the sub-epidermal tissues of the root [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, the GAs/DELLAs have a role in the production of trichomes (leaf hairs) in *A. thaliana* [CHIEN & SUSSEX, 1996; TRAW & BERGELSON, 2003] and participate in the organization of the cytoskeleton of microtubules (MT) [LOCASCIO & al. 2013], which is essential for trichome and root hair growth, for establishing root cell identity and shape, and for plant cell expansion and division [BAO & al. 2001].

Because auxin, ethylene, abscisic acid, nitric oxide, brassinosteroids, cytokinins and strigolactones regulate the root hair patterning [CAO & al. 1999; VAN HENGEL & al. 2004; LOMBARDO & al. 2006; KAPPUSAMY & al. 2009; NIU & al. 2011], and because changes in the levels of these phytohormones are correlated to alterations in the root cell size/number and the root architecture in response to nutritional stresses, such as low availability of P, B or Fe in the soil (e.g. swelling of root cortex cells, induction of extra cells at the root cortex, and production of lateral roots (LR)) [SCHMIDT & al. 2000; YANG & al. 2007; MARTÍN-REJANO & al. 2011], this study wanted to determine whether supra-physiological levels of GAs/DELLAs might also have an effect on the size/number of root cells and on the production of LR in root tips of *A. thaliana* seedlings. To this aim, the size and/or number of root tip cells

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

were examined in Col (0) seedlings grown for 5 days under excessive levels of GAs/DELLAs, as well as in GAs (*QD*, *5X*, *GID1b-ox*) or DELLAs (*gai-1*, *HSp::gai-1*, *pGAI::gai-1:GR*, *SCR::gai-1:GR*)-overproducing mutants. Moreover, the size of root tip cells was examined in 5-day-old mutant seedlings resulting from expressing the *gai-1* (GA-insensitive) DELLA allele in different tissues of the root (UAS (GAL4-UPSTREAM ACTIVATION SEQUENCE) expression directed system lines; Dr. JIM HASSELHOFF'S laboratory). On the other hand, the presence of LR was also analysed in root tips of *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Results of this study suggested that the GAs/DELLAs might have a role in regulating the size, number and organization of root cells, as well as the root architecture, in root tips of *A. thaliana* seedlings.

Material and methods

Plant Material and Growth Conditions

Arabidopsis thaliana Col (0) seeds were sterilized (70% Ethanol (v/v) and 0.01% Triton X-100 (v/v)), sown on half-strength MS medium plates (0.8% (w/v) agar and 1% (w/v) sucrose), stratified for 3-4 days (4 °C, darkness), germinated, and grown vertically (22 °C; 5-7 days) under continuous white light (Percival growth chamber E-30B) (<http://www.percival-scientific.com>) as described by LEE & SCHIEFELBEIN (1999).

Hormone and Chemical treatments

Stock solutions of paclobutrazol (PAC, 10 mM in acetone 100% (v/v)), GA₄ (1 mM in 100% ethanol (v/v)) or GA₃ (50 mM in 100% ethanol (v/v)) were conveniently diluted and added to MS agar medium or water (in the case of liquid incubation experiments) to obtain a final concentration of 0.5 μM PAC, 1 μM GA₄ and 30 μM GA₃.

Mutant Lines

In a previous study [MCCARTHY-SUÁREZ, unpublished manuscript], the spatial gene expression of the root non-hair epidermal cell fate marker *GL2* was studied in root tips of *A. thaliana* seedlings by using the *GL2pro::GUS* mutant line as well as those derived from crossing lines harbouring constitutively excessive levels of GAs/DELLAs with the *GL2pro::GUS* line (*Ler* x *GL2pro::GUS* background). In the present study, the effect of transient increases in the levels of expression of the *gai-1* (GA-insensitive-1) DELLA allele on the size and number of root tip cells in *A. thaliana* seedlings was examined by using the *gai-1* mutant lines of heat-shock inducible *HSp::gai-1* (which over-expresses *gai-1* upon heat shock) and of dexamethasone (DEXA)-inducible *pGAI::gai-1:GR* and *SCR::gai-1:GR* (with glucocorticoid-binding domain). The *HSp::gai-1* mutant seedlings were grown at 37 °C for 4 h (heat-shock) and then at 22 °C for 2 h (recovery period), whereas the *pGAI::gai-1:GR* and *SCR::gai-1:GR* mutant seedlings were incubated in 0.2, 1.2 or 10 μM DEXA for a minimum of 6h. Root cell size and number was also studied in mutants with excessive levels of GAs/DELLAs (*gai-1*, *QD* (*quadruple DELLA mutant*), *5X* (*quintuple DELLA mutant*) and *GID1b-ox* (which over-expresses the GA receptor *GID1b* (GIBBERELLIN INSENSITIVE DWARF1)), in mutants over-expressing *gai-1* in different tissues of the root (*ML1::gai-1* (epidermis) and UAS expression directed system (GAL4-UPSTREAM ACTIVATION SEQUENCE) lines: *UAS::gai-1* x C24 (control, background); *UAS::gai-1* x J0951 (epidermis of the meristematic zone (MZ)); *UAS::gai-1* x J2812 (MZ epidermis and cortex); *UAS::gai-1* x N9142 (cortex of elongation zone (EZ)); *UAS::gai-1* x M0018 (MZ cortex and endodermis);

UAS::gai-1 x J0571 (MZ cortex and endodermis); *UAS::gai-1* x Q2393 (all tissues but the endodermis); *UAS::gai-1* x Q2500 (MZ endodermis/pericycle); *UAS::gai-1* x J0121 (EZ pericycle); *UAS::gai-1* x J0631 (all tissues of the EZ); *UAS::gai-1* x J3281 (vessels)), and in the *35S::CPC* x *GL2pro::GUS* and *scm* (scrambled) x *GL2pro::GUS* mutants.

GUS activity assay

GUS (β -glucuronidase) staining of the *GL2pro::GUS* reporter line was performed as described by FRIGERIO & al. (2006), but using 8 mM instead of 2 mM potassium ferro/ferricyanide and incubating the seedlings (15 min to 2 h) in the reaction mixture at 4 °C instead of 37 °C.

Microscopy

Cell organization at the root tip was studied on ultra-thin cross sections of plastic resin-embedded roots as described by Dr. SCHIEFELBEIN Protocols (<http://www.mcdb.lsa.umich.edu/labs/schiefel/protocols.html>). Seedlings were included in 1% agarose in 0.1 M sodium phosphate buffer, pH 6.8, and stained for GUS activity. Root-containing blocks were then cut, fixed with 4% para-formaldehyde in PBS, dehydrated in ethanol series (15%, 30%, 50%, 75%, 95% and 100%, 1 h each), kept in 100% ethanol overnight, incubated in Technovit ® 7100 infiltration solution for 2 days, inserted in gelatine capsules, and embedded for 9 days in Technovit ® 7100 plastic resin (Heraeus Kultzer, Germany). Ultramicrotome (Ultracut E, Reichert Jung, Germany) cross sections of resin-embedded roots were then stained with 0.06% (w/v) toluidine blue and observed under a Nikon Eclipse E600 microscope. The GFP expression in HASELHOFF's crossed lines was visualized by using a Leica Confocal Microscope (Excitation: 488 nm; Detection: 500-530 nm band-path filter for GFP).

Results

Excessive levels of GAs/DELLAs modified the size and number of root tip cells in seedlings of *A. thaliana*

Apart from altering the patterning, morphology and abundance of root hairs [MCCARTHY-SUÁREZ, unpublished manuscript], high levels of GAs/DELLAs also modified the size and number of root tip cells in seedlings of *A. thaliana* (Figures 1-3; Tables 1-3). While excessive levels of GAs, excepting the epidermal cells, usually narrowed the root cells, excessive levels of DELLAs frequently widened, shortened and twisted the root epidermal, cortical, endodermal and pericycle cells, resulting in wider roots (Figures 1-3; Tables 1-3). Moreover, in the *HSp::gai-1* and *SCR::gai-1:GR* mutants, the widening and shortening of the root epidermal, cortical, endodermal and pericycle cells observed at 24 h after heat-shock (37 °C, 4 h) or after growth in DEXA (10 μ M), respectively (Figure 3), was accompanied, as previously reported [MCCARTHY-SUÁREZ, unpublished manuscript], by an alteration in the spatial expression of *GL2* and in the distribution of root hairs. Similar changes were observed when *gai-1* was over-expressed at the subepidermal tissues of the root (Figures 1, 2 and 4; Table 1). However, when *gai-1* was over-expressed at the root epidermis, no apparent changes occurred in the size of the root epidermal, cortical, endodermal or pericyclic cells (Figures 1, 2 and 4; Table 1), in the patterning of *GL2* gene expression or in the root hair distribution [MCCARTHY-SUÁREZ, unpublished manuscript], what suggested that the changes in root cell size that took place when *gai-1* was over-expressed at the sub-epidermal tissues of the root might have been

connected to the alterations in the root hair patterning induced by excessive levels of GAs/DELLAs.

Growth of *A. thaliana* seedlings for 5 days under excessive levels of GAs, in contrast, caused the narrowing of the root cortical cells, an effect that was corroborated in the *QD*, *5X* and *GID1b ox* mutants (Figure 2; Table 1). Nevertheless, excessive levels of GAs also seemed to slightly increase the relative width of the root epidermal cells (Figures 1, 5 and 6). Frequently, under GA_3 (30 μ M) or GA_4 (1 μ M), changes of cell fate at the root epidermis coincided with changes in the width of the epidermal cells and/or with changes in the size of the root cortical and endodermal cells (Figure 3; Tables 1 and 2).

In fact, estimations of tissue depth in root tips of *A. thaliana* seedlings uncovered the swelling of the root epidermal, cortical, endodermal and pericyclic cells under high levels of DELLAs (PAC) and their slight thinning under high levels of GAs (*5X* mutant) (Table 4). Moreover, growth of seedlings of the *scm x GL2pro::GUS* mutant in PAC (0.5 μ M) for 5d caused the radial swelling of the epidermal, cortical and endodermal and pericyclic cells of the root tip (Figure 6). On the other hand, in seedlings of the *35S::CPC x GL2pro::GUS* mutant, changes of cell fate at the root epidermis were accompanied by changes in the width of the underlying root epidermal and cortical cells (Figure 3).

Interestingly, cell size changes in root tips under excessive levels of GAs/DELLAs were often accompanied by the presence of multinucleated cells at the epidermis, cortex, endodermis and pericycle of the root (Figure 6).

Excessive levels of GAs/DELLAs also modified the radial cell organization in root tips of *A. thaliana* seedlings (Figure 5). Treatments with PAC (5d, 7d) frequently increased the number of cells at the epidermis, cortex, endodermis and pericycle of the root (Figures 5 and 6; Table 5) and induced anticlinal/diagonal cell divisions at the root epidermis (T-clones) (Figure 3) as well as periclinal cell divisions at the root cortex and endodermis (middle cortex (MC)) (Figure 6). Furthermore, growth of the *scm x GL2pro::GUS* mutant for 5 days in PAC induced the proliferation of the root epidermal, cortical and endodermal cells and the formation of a MC (Figure 6). Treatments with excessive levels of GAs, in turn, sometimes increased the number of cortical and pericycle cells, but not of epidermal cells, in the radial dimension of the root, and induced epidermal T-clones and a MC (Figures 3, 5 and 6; Table 5). Moreover, the number of root epidermal cells decreased in the *5X* mutant (Figure 5; Table 5), which, in part, might have explained the lower abundance of root hairs per root radial section in this mutant [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, in root cross sections, frequently only one epidermal cell was seen at the Atrichoblast (non-hair) position under high levels of GAs, whereas up to four cells could be seen under high levels of DELLAs (Figure 5), in tune with the reduced number of epidermal cells in the *5X* mutant and the increased number of epidermal cells under PAC (Table 5). Furthermore, given that root non-hair cells lay over just one cortical cell, then, the observed increase in the width of the cortex cells under high levels of DELLAs (Figures 2, 4 and 5; Table 1) might have accounted for the higher percentage of epidermal cells at the Atrichoblast position, as well as the lower percentage of epidermal cells at the Trichoblast position, detected per root radial section under this treatment [MCCARTHY-SUÁREZ, unpublished manuscript]. Conversely, the decrease in the width of the cortex cells seen under high levels of GAs (*5X* mutant) (Figure 2; Table 1) might have explained the lower percentage of epidermal cells at the Atrichoblast position, and the higher percentage of epidermal cells at the Trichoblast position, found per root radial section under this treatment [MCCARTHY-SUÁREZ, unpublished manuscript]. Nevertheless, considering that the average number of epidermal cells per root radial section increased under high levels of DELLAs (PAC,

gai-1) and decreased under high levels of GAs (5X mutant) [MCCARTHY-SUÁREZ, unpublished manuscript], then, the predicted number of epidermal cells at the Trichoblast position did not change under excessive levels of GAs/DELLAs in 5 day-old *A. thaliana* seedlings [MCCARTHY-SUÁREZ, unpublished manuscript].

The root diameter at the MZ, in addition, increased by 40% under excessive levels of DELLAs (Table 3), in tune with the increased number of cells at the root epidermis, cortex, endodermis and pericycle (Figure 5; Table 5) and the wider and/or deeper cells at the root cortex, endodermis and pericycle (Figures 5 and 6; Tables 1 and 4). Conversely, under excessive levels of GAs, the root diameter decreased (Table 3), in accordance with the lower number of cells at the root epidermis (5X mutant) (Table 5), and the narrower and shallower cells at the root cortex and endodermis (Figures 2, 5 and 6; Tables 1 and 4). Nevertheless, at the MZ-EZ transition zone, the root also seemed to swell under excessive levels of GAs, and there was variability of cell sizes [MCCARTHY-SUÁREZ, unpublished manuscript] (Figure 6), maybe due to the swelling of the epidermal cells and to the deepening of the pericycle cells (Figure 1; Table 4). In fact, GA₃ treatments increase the ratio of (xylem/whole root) area [WANG & al. 2015].

Excessive levels of GAs / DELLAs modified the outgrowth of lateral roots in root tips of *A. thaliana* seedlings

Excessive levels of DELLAs also induced the outburst of LR near the root tip in *A. thaliana* seedlings, whereas excessive levels of GAs inhibited it (Figure 7).

Table 1. Average length and width of root cortical cells in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Analyses performed on micrographs of root cortex cells, 20X. (*) At 48h after a 4h-heat-shock experiment.

	Root cortical cell			
	N° Cells analysed	Length (µm)	N° Cells analysed	Width (µm)
Col (0) (MS)	25	171 ± 28 (100 %)	66	31 ± 3 (100 %)
PAC (0.5 µM)	23	153 ± 34 (89 %)	60	43 ± 3 (142 %)
GA ₄ (1 µM)	26	209 ± 52 (122 %)	26	27 ± 2 (87 %)
PAC (0.5 µM) + GA ₄ (1 µM)	24	173 ± 49 (101 %)	28	19 ± 2 (61 %)
<i>Ler</i>	28	188 ± 29 (100 %)	18	30 ± 4 (100 %)
<i>ML1::gai-1</i>	12	171 ± 33 (91 %)	28	30 ± 3 (97 %)
<i>gai-1</i>	71	138 ± 41 (73 %)	72	37 ± 7 (123 %)
<i>QD</i>	33	180 ± 34 (96 %)	53	27 ± 3 (90 %)
<i>5X</i>	15	187 ± 34 (99 %)	30	27 ± 4 (90 %)
<i>GID1b-ox</i>	23	210 ± 54 (112%)	34	24 ± 4 (80 %)
<i>Hsp::gai-1 x GL2pro::GUS</i> (22 °C, 4 h)	48	184 ± 39 (100 %)	47	27 ± 4 (100 %)
<i>Hsp::gai-1 x GL2pro::GUS</i> (37 °C, 4 h)	38	158 ± 52 (86 %)	33	31 ± 4 (115 %)
<i>Hsp::gai-1 x GL2pro::GUS</i> (22 °C) 48 h*	23	195 ± 43 (100 %)	34	29 ± 3 (100%)
<i>Hsp::gai-1 x GL2pro::GUS</i> (37 °C) 48 h*	36	61 ± 19 (31 %)	44	46 ± 9 (159 %)
<i>pGAI::gai-1:GR</i> (MS)	20	180 ± 56 (100 %)	20	34 ± 4 (100 %)
<i>pGAI::gai-1:GR</i> (10 µM DEXA)	29	129 ± 31 (72 %)	29	46 ± 5 (135 %)
<i>SCR::gai-1:GR</i> (MS)	77	140 ± 51 (100 %)	77	34 ± 5 (100%)
<i>SCR::gai-1:GR</i> (10 µM DEXA)	64	70 ± 32 (50 %)	63	53 ± 10 (156 %)
<i>UAS::gai-1 x C24</i> (control)	46	154 ± 49 (100 %)	52	28 ± 4 (100 %)
<i>UAS::gai-1 x J0951</i> (epidermis)	54	155 ± 42 (101%)	72	31 ± 4 (111 %)
<i>UAS::gai-1 x J2812</i> (epi + cortex)	79	116 ± 48 (75 %)	91	35 ± 6 (125 %)

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

<i>UAS::gai-1 x J0571</i> (cortex + endo)	74	71 ± 18 (46 %)	74	48 ± 8 (171 %)
<i>UAS::gai-1 x M0018</i> (cortex + endo)	49	89 ± 31 (58 %)	46	46 ± 9 (164 %)
<i>UAS::gai-1 x Q2500</i> (MZ endo/pericycle)	79	51 ± 13 (33 %)	85	41 ± 8 (146 %)
<i>UAS::gai-1 x Q2393</i> (all but endo)	23	174 ± 59 (113 %)	32	32 ± 5 (114 %)
<i>UAS::gai-1 x J0631</i> (elong. tissues)	73	46 ± 8 (30 %)	67	37 ± 6 (132 %)
<i>UAS::gai-1 x J0121</i> (EZ pericycle)	16	177 ± 45 (115 %)	19	27 ± 3 (96 %)

Table 2. Cortical and endodermal cell area in cross sections of root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Measurements of transversal cell length and width were performed on micrographs of cross sections of resin-embedded roots (40X). Area was calculated by multiplying cell length (µm) by cell width (µm).

	N° cross sections analysed	Cortical cell area (µm ²)	N° cross sections analysed	Endodermal cell area (µm ²)
Col (0) (MS)	84	507 ± 7 (100 %)	82	169 ± 3 (100 %)
PAC (0.5 µM)	105	737 ± 17 (145 %)	82	215 ± 7 (127 %)
GA ₄ (1 µM)	98	375 ± 13 (74 %)	81	157 ± 5 (93 %)
<i>Ler</i>	8	509 ± 9 (100 %)	24	172 ± 4 (100 %)
<i>gai-1</i>	100	464 ± 5 (91 %)	89	186 ± 4 (108 %)
<i>QD</i>	16	264 ± 6 (52 %)	23	138 ± 4 (80 %)
<i>5X</i>	7	270 ± 2 (53 %)	51	143 ± 3 (83 %)

Table 3. Average width of the root MZ in root tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Measurements of root width were performed on micrographs of whole root tips (10X, longitudinal view) or cross sections of resin-embedded roots (40X).

Root width (µm)				
	N° roots analysed	Whole root tips	N° cross sections analysed	Root cross sections
Col (0) (MS)	12	115 ± 4 (100 %)	84	144 ± 10 (100 %)
PAC (0.5 µM)	19	161 ± 12 (140 %)	107	164 ± 14 (114 %)
GA ₄ (1 µM)	19	99 ± 12 (86 %)	96	128 ± 18 (89 %)
<i>Ler</i>	10	122 ± 5 (100 %)	8	145 ± 4 (100 %)
<i>gai-1</i>	18	125 ± 18 (102 %)	100	148 ± 13 (102 %)
<i>QD</i>	16	107 ± 18 (88 %)	41	129 ± 7 (89 %)
<i>5X</i>	5	111 ± 0 (91 %)	50	132 ± 8 (91 %)


Table 4. Estimated tissue depth in roots tips of 5-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. Measurements of tissue diameter (μm) were performed on micrographs of cross sections of resin-embedded roots (MZ to early EZ; 40X). Estimated epidermal depth = [Root diameter (data from table 3) – (Cortex-Endodermis-Pericycle-Vessels diameter)]/2. Estimated cortical depth = [(Cortex-Endodermis-Pericycle-Vessels diameter) – (Endodermis-Pericycle-Vessels diameter)]/2. Estimated endodermal depth = [(Endodermis-Pericycle-Vessels diameter) – (Pericycle-Vessels diameter)]/2. Estimated pericycle depth = [(Pericycle-Vessels diameter) – (Vessels diameter)]/2. Endo: Endodermis; Peri: Pericycle. Number of cross sections analysed: Control (27-29), PAC (46-49), GA₄ (74-78), *Ler* (21-22), *gai-1* (36), *QD* (34-36), *5X* (47-48).

	Col (0) (MS)	PAC (0.5 μM)	GA ₄ (1 μM)	<i>Ler</i>	<i>gai-1</i>	<i>QD</i>	<i>5X</i>
Cortex-Endo-Peri-Vessels diameter	95 \pm 6 (100%)	110 \pm 11 (116%)	83 \pm 12 (87%)	90 \pm 7 (100%)	91 \pm 5 (101%)	81 \pm 5 (90%)	82 \pm 4 (91%)
Endo-Peri-Vessels diameter	59 \pm 6 (100%)	70 \pm 7 (119%)	58 \pm 7 (98%)	60 \pm 4 (100%)	59 \pm 3 (98%)	59 \pm 6 (98%)	55 \pm 2 (92%)
Peri-Vessels diameter	45 \pm 5 (100%)	54 \pm 5 (120%)	45 \pm 6 (100%)	45 \pm 3 (100%)	44 \pm 2 (98%)	45 \pm 5 (100%)	40 \pm 2 (89%)
Vessels diameter	35 \pm 3 (100%)	40 \pm 3 (114%)	32 \pm 5 (91%)	33 \pm 3 (100%)	34 \pm 2 (103%)	33 \pm 4 (100%)	30 \pm 2 (91%)
Estimated Epidermal depth	25 (100%)	27 (108%)	22 (88%)	27 (100%)	28 (104%)	24 (89%)	25 (93%)
Estimated Cortex depth	18 (100%)	20 (111%)	13 (72%)	15 (100%)	16 (107%)	11 (73%)	13 (87%)
Estimated Endodermis depth	7 (100%)	8 (114%)	6 (86%)	8 (100%)	7 (88%)	7 (88%)	7 (88%)
Estimated Pericycle depth	5 (100%)	7 (140%)	6 (120%)	6 (100%)	5 (83%)	6 (100%)	5 (83%)

Table 5. Radial cell organization in roots of 5 or 7-day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLA. Analyses performed on micrographs of cross sections of resin embedded-roots (40X).

	N° root cross sections analysed	Epidermis	Cortex	Endodermis	Pericycle
Control (5d)	75	20–25	8	8–10	13–15
PAC (5d)	56	25–34	8–9	9–13	14–18
GA₄ (5d)	56	17–25	8–9	8–10	12–16
Control (7d)	14	24–27	8	8–10	14–15
PAC (7d)	18	26–33	8–10	11–13	14–18
GA₄ (7d)	45	23–25	8–9	8–10	14–16
<i>Ler</i> (5d)	13	18–24	8	8	13–14
<i>gai-1</i> (5d)	57	19–25	8	8–9	12–14
<i>QD</i> (5d)	24	22–24	8–11	8–11	14–18
<i>5X</i> (5d)	54	18–21	8–9	7–8	12–16

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...






Figure 2

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...




Figure 3




Figure 4




Figure 5




Figure 6

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...



Figure 7

Figure legends

Figure 1. Excessive levels of GAs/DELLAs altered the size of root epidermal cells in root tips of 5-day-old *A. thaliana* seedlings. **A)** Col(0) (MS); **B)** Col(0) (0.5 μ M PAC); **C)** Col(0) (30 μ M GA3); **D)** Col(0) (1 μ M GA4); **E)** *Ler*; **F)** *gai-1*; **G)** *QD*; **H)** *5X*; **I)** *GID1b-ox* (MS); **J)** *HSp::gai-1* (22 $^{\circ}$ C, 4 h); **K)** *HSp::gai-1* (immediately after heat shock (37 $^{\circ}$ C, 4 h); **L)** *HSp::gai-1* (24 h after heat-shock (37 $^{\circ}$ C, 4 h); **M)** *HSp::gai-1* (48 h after heat-shock (37 $^{\circ}$ C, 4 h); **N)** *pGAI::gai-1:GR* (MS); **O)** *pGAI::gai-1:GR* (10 μ M DEXA); **P)** *SCR::gai-1:GR* (MS, leaky line); **Q)** *SCR::gai-1:GR* (10 μ M DEXA); **R)** *UAS::gai-1* x C24; **S)** *UAS::gai-1* x J0951; **T)** *UAS::gai-1* x J2812; **U)** *UAS::gai-1* x J0571; **V)** *UAS::gai-1* x Q2500; **W)** *UAS::gai-1* x J0121; **X)** *UAS::gai-1* x J0631; **Y)** *UAS::gai-1* x J3281. Magnification: 20X. The scale bar represents 100 μ m. Propidium iodide staining.

Figure 2. Excessive levels of GAs/DELLAs altered the size of root cortical cells in root tips of 5-day-old *A. thaliana* seedlings. **A)** Col(0) (MS); **B)** Col(0) (0.5 μ M PAC); **C)** Col(0) (1 μ M GA4); **D)** *Ler*; **E)** *gai-1*; **F)** *QD*; **G)** *5X*; **H)** *GID1b-ox*; **I)** *pGAI::gai-1:GR* (3d in MS); **J)** *pGAI::gai-1:GR* (3d in 10 μ M DEXA); **K)** *SCR::gai-1:GR* (3d in MS); **L)** *SCR::gai-1:GR* (3d in 10 μ M DEXA); **M)** *HSp::gai-1* x *GL2pro::GUS* (24h after 22 $^{\circ}$ C for 4 h); **N)** *HSp::gai-1* x *GL2pro::GUS* (24h after heat-shock (37 $^{\circ}$ C for 4 h)); **O)** *GL2pro::GUS* (heat-shock control) (24 h after 22 $^{\circ}$ C for 4 h); **P)** *GL2pro::GUS* (heat-shock control) (24h after heat shock (37 $^{\circ}$ C for 4 h)); **Q)** *UAS::gai-1* x C24; **R)** *ML1::gai-1*; **S)** *UAS::gai-1* x J2812; **T)** *UAS::gai-1* x M0018; **U)** *UAS::gai-1* x Q2500; **V)** *UAS::gai-1* x N9142; **W)** *UAS::gai-1* x J0121; **X)** *UAS::gai-1* x 0631; **Y)** *UAS::gai-1* x J3281. Magnification: 20X. The scale bar represents 100 μ m. Propidium iodide staining.

Figure 3. Excessive levels of GAs/DELLAs induced cell size changes and T-clones at the epidermis, cortex and endodermis of root tips of 5-day-old *A. thaliana* seedlings. **A)** *GL2pro::GUS* (30 μ M GA3) (all layers): expansion of an epidermal cell and narrowing of a cortical cell coincide with a change in epidermal cell fate, 40X; **B)** *35S::CPC* x *GL2pro::GUS* (epidermis): A change in epidermal cell size coincides with a change in epidermal cell fate, 40X; **C)** *35S::CPC* x *GL2pro::GUS* (cortex): A change in the width of a cortical cell coincides with a change in epidermal cell fate, 40X; **D)** *35S::CPC* x *GL2pro::GUS* (all layers): epidermis and cortex vary in cell size, 40X; **E)** *HSp::gai-1* x *GL2pro::GUS* (48 h after 22 $^{\circ}$ C for 4 h) (all layers), 20X; **F)** *HSp::gai-1* x *GL2pro::GUS* (48h after heat-shock (37 $^{\circ}$ C, 4 h) (all layers): swelling of the root epidermal, cortical, endodermal and pericyclic cells, 20X; **G)** *SCR::gai-1:GR* (MS) (all layers), 20X; **H)** *SCR::gai-1:GR* x *GL2pro::GUS* (24 h in 10 μ M DEXA) (all layers): swelling of the root cortex, endodermis and pericycle, 20X; **I)** Epidermal T-clones in PAC (0.5 μ M), 40X; **J)** Epidermal T-clones in GA3 (30 μ M), 40X; **K)** *SCR::gai-1:GR* (0.2 μ M DEXA): epidermal T-clones, 20X; **L)** *SCR::gai-1:GR* (1.2 μ M DEXA): periclinal cell division at the root cortex, 20X; **M)** *SCR::gai-1:GR* (MS) (all layers): arrow on endodermis, 40X; **N)** *SCR::gai-1:GR* (10 μ M DEXA) (all layers): periclinal cell divisions at the endodermis, 40X; **O)** *SCR::gai-1:GR* (10 μ M DEXA) (all layers): periclinal cell divisions at the endodermis, 40X. The scale bar represents 100 μ m (20X) or 50 μ m (40X). Propidium iodide or GUS staining.

Figure 4. Over-expression of *gai-1* in different tissues of the root modified the cell size in root tips of 5-day-old *A. thaliana* seedlings. **A)** *UAS::gai-1* x C24 (Background); **B)** *UAS::gai-1* x J0951 (epidermis of the MZ); **C)** *UAS::gai-1* x J2812 (epidermis and cortex of the MZ); **D)** *UAS::gai-1* x N9142 (cortex of EZ); **E)** *UAS::gai-1* x J0571 (cortex and endodermis of the MZ); **F)** *UAS::gai-1* x Q2500 (endodermis/pericycle of MZ); **G)** *UAS::gai-1* x J0121 (pericycle of EZ); **H)** *UAS::gai-1* x Q2393 (all tissues but the endodermis); **I)** *UAS::gai-1* x J0631 (elongating tissues); **J)** *UAS::gai-1* x J3281 (vessels). Magnification: 40X. The scale bar represents 50 μ m. Propidium iodide staining.

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

Figure 5. Radial cell organization in root tips of 5 or 7 day-old *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. **A)** Col(0) (MS, 5d); **B)** Col(0) (0.5 μ M PAC, 5d); **C)** Col(0) (1 μ M GA4, 5d); **D)** Col(0) (MS, 7d); **E)** Col(0) (0.5 μ M PAC, 7d); **F)** Col(0) (1 μ M GA4, 7d); **G)** *Ler* (5d); **H)** *gai-1* (5d); **I)** *QD* (5d); **J)** *5X* (5d). Magnification: 40X. The scale bar represents 50 μ m. Toluidine blue staining.

Figure 6. Excessive levels of GAs/DELLAs induced multinucleated cells, a middle cortex (MC) and extra cortical cells in root tips of *A. thaliana* seedlings. **A)** PAC (0.5 μ M, 5d): Epidermal multinucleated cell; **B)** PAC (0.5 μ M, 5d): Cortical and endodermal multinucleated cells; **C)** PAC (0.5 μ M, 5d): Pericyclic multinucleated cell; **D)** GA4 (1 μ M, 5d): Epidermal and cortical multinucleated cells; **E)** PAC (0.5 μ M, 5d): MC; **F)** PAC (0.5 μ M, 7d): MC (arrow) and 10 cortical cells; **G)** GA4 (1 μ M, 5d): MC (arrow) and 9 cortical cells; **H)** *scm* x *GL2pro::GUS* (MS, 5d); **I)** *scm* x *GL2pro::GUS* (0.5 μ M PAC, 5d): MC (arrow) and 9 cortical cells. Magnification: 40X. The scale bar represents 50 μ m. Toluidine blue staining.

Figure 7. Over-expression of *gai-1* induced an early outburst of lateral roots in root tips of *A. thaliana* seedlings. **A)** *SCR::gai-1:GR* (MS, 5d) (Leaky line), 4X; **B)** *SCR::gai-1:GR* (10 μ M DEXA, 5d), 4X; **C)** *UAS::gai-1* x C24 (control, 5d), 4X; **D)** *UAS::gai-1* x J2812 (5d), 20X; **E)** *UAS::gai-1* x M0018 (5d), 4X; **F)** *UAS::gai-1* x J0571 (5d), 4X; **G)** *UAS::gai-1* x Q2500 (5d), 4X; **H)** *UAS::gai-1* x N9142 (5d), 20X; **I)** *UAS::gai-1* x J3281 (8d) (aborted primary root), 4X. The scale bar represents 100 μ m (20X) or 500 μ m (4X).

Discussion

The GAs/DELLAs might regulate the size and number of root tip cells in seedlings of *A. thaliana*.

Root cell size: Connections to the root hair patterning and abundance

Apart from altering the patterning, morphology and abundance of root hairs [MCCARTHY-SUÁREZ, unpublished manuscript], results of the present study suggest that excessive levels of GAs/DELLAs also modified the size of root tip cells in seedlings of *A. thaliana*. While excessive levels of DELLAs frequently shortened and widened the epidermal, cortical, endodermal and pericyclic cells of the root, what resulted in wider root tips, excessive levels of GAs, with the exception of the epidermal cells, often narrowed them. Thus, because root hair cells are shorter than the root non-hair cells [SALAZAR-HENAO & al. 2016], then, the inhibition of epidermal cell elongation that occurred when *gai-1* was over-expressed in tissues placed underneath the MZ epidermis (*J2812* >> *gai-1*, *J0571* >> *gai-1*, *M0018* >> *gai-1*, *Q2500* >> *gai-1* or *Q2393* >> *gai-1* lines) or in all tissues of the EZ (*J0631* >> *gai-1* line) might have contributed, in part, to the appearance of ectopic root hairs, and, therefore, to the higher density of root hairs observed near the root tip under excessive levels of DELLAs [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, *Arabidopsis* increases root hair density by decreasing the length of root epidermal cells, as shown under P deficiency [JIANG & al. 2007; PÉRET & al. 2011; SALAZAR-HENAO & al. 2016; JANES & al. 2018]. Alternatively, given that GAs promote the elongation of root epidermal cells, accumulate at the endodermis of the root EZ, and affect the expansion of root EZ cells by destabilizing DELLAs and inducing expansin genes [UBEDA-TOMÁS & al. 2008, 2009; GOU & al. 2010; BAHIN & al. 2011; SHANI & al. 2013], then, the extra elongation of root epidermal cells which is known to occur under high levels of GAs [BAND & al. 2012] might have contributed, in turn, to the lower density of root hairs that was observed at the root tip under this treatment [MCCARTHY-

SUÁREZ, unpublished manuscript]. In fact, in WT, cell expansion at the root EZ is strictly polar and is not accompanied by an increase in the root diameter [BAO & al. 2001].

Interestingly, epidermal patterning genes instruct epidermal cell size [LÖFKE & al. 2013]. In turn, variations in the expression of HDA 19 (histone deacetylase 19), which controls epidermal cell elongation, affect root cell elongation and, thus, root hair density [CHEN & al. 2015; SALAZAR-HENAO & al. 2016]. In this study, however, neither the patterning or abundance of root hairs [MCCARTHY-SUÁREZ, unpublished manuscript] nor the epidermal cell length seemed to suffer any alterations when *gai-1* was over-expressed at the root epidermis (Figure 4), what suggests that the changes of cell size induced by excessive levels of GAs/DELLAs in roots of *A. thaliana* seedlings might have been orchestrated from the sub-epidermal tissues of the root. Furthermore, WILD & al. (2016) showed that expressing *gai-1* at the root epidermis did not affect the root length.

Results of this study also seem to point at a link between cell size and cell fate, because expressing *gai-1* at the elongating tissues of the root (J0631 >> *gai-1*) caused the shortening and widening of the majority of the root cells (Figures 1, 2 and 4) along with a hairy phenotype similar to that of the *wer* mutant [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, the DELLAs inhibit the elongation of root EZ cells and primary root [ALONSO-RAMÍREZ & al. 2009; UBEDA-TOMÁS & al. 2009; LEE & al. 2012] and down-regulate PIF4, a phytochrome-interacting factor which induces cell elongation genes [ACHARD & GENSHIK, 2009]. Deficiencies in B, P or Fe, which increase the levels of DELLAs at the root MZ and induce ectopic root hairs, also reduce the primary root length [MARTÍN-REJANO & al. 2011; PÉRET & al. 2011; WILD & al. 2016]. Furthermore, it has been suggested that production of ectopic root hairs in ectopic root hair 2 (*erh2*) occurs at late stages of root development, correlated with cell expansion. However, it has also been suggested the independence of root hair initiation from cell expansion, as *erh3* acts as soon as cell fate specification [SCHNEIDER & al. 1997]. Particularly, ERH3 is required for the stable fixation of positional signals at the cell wall (CW) for cell fate specification. Moreover, ERH3 codifies a MT-severing p60 katanin protein and has a role in CW biosynthesis [WEBB & al. 2002]. On the other hand, given that the expression of cell identity markers is altered in *erh3*, it has been suggested that MT are directly active in the specification of root cell identity, and that MT disruption in *erh3* results in the development of defective identities [WEBB & al. 2002]. In fact, in several animal systems, MT are involved in the specification of cell identity and polarity [WEBB & al. 2002].

Shortening and radial expansion of root cortical and endodermal cells has also been reported in the *cobra*, *pom-1* (both cellulose deficient), *shoebbox* (GA biosynthesis-impaired), *dgl1* (GA-insensitive), TUA6/AS (α -tubulin-deficient), *erh*, *sabre*, *PLD* (phospholipase D), *wer*, *scm* and *jkd* mutants, in plants treated with umbelliferone (a cellulose biosynthesis inhibitor), MT-breaking drugs or 1-butanol (an inhibitor of PLD), and in plants stressed by salinity, gamma irradiation or mineral deficiency (Fe, P) [JANKAY & MULLER, 1976; SCHIEFELBEIN & al. 1997; SCHNEIDER & al. 1997; BAO & al. 2001; MA & al. 2001; SCHERES & al. 2002; GARDINER & al. 2003; NAGATA & al. 2004; KOMORISONO & al. 2005; WELCH & al. 2007; DINNENY & al. 2008; PIETRA & al. 2015; JANES & al. 2018]. Cortical cell expansion as a result of excessive DELLAs has also been described by BENFEY & al. [1993]. Interestingly, PAC induces the expression of expansin genes at the root cortex (At4g21280 (+2,002); Arex data), being *EXPANSIN 7* a specific marker of root hair cells [OHASHI & al. 2003; GENDRE & al. 2019]. In addition, it has been suggested that the DELLA GAI might have a role in the expansion of endodermal cells in *Arabidopsis* primary roots, and that the expansion of endodermal cells determines the elongation of whole roots [UBEDA-

TOMÁS & al. 2009; ZHANG & al. 2014). On the other hand, the GAs control the size of the root apical meristem (RAM) in *Arabidopsis* by affecting cortical cell expansion [NELISSEN & al. 2012; FONOUNI-FARDE & al. 2019]. In fact, for different accessions of *Arabidopsis*, there is a correlation between cortex cell length and the length of the root MZ [ZHANG & al. 2014]. Furthermore, when *gai-1* is over-expressed at the root endodermis (the most important tissue for GA-dependent root growth), the cessation of anisotropic cell growth expands radially the cortical cells and causes the outward protrusion of epidermal cells [UBEDA-TOMÁS & al. 2008]. With this regard, it is known that the endodermis of the EZ regulates nutrient uptake [PÉRET & al. 2011; SHANI & al. 2013; CUI, 2015], whereas the cortex participates in the root response to P deficiency [SHIN & al. 2005].

Therefore, the shortening and widening of the root epidermal, cortical, endodermal and pericycle cells induced by excessive levels of DELLAs in seedlings of *A. thaliana* might have explained the radial expansion of the root tips observed. Interestingly, treatment with PAC also increased the root diameter in carrot [WANG & al. 2015]. In fact, in this study, root tips became thinner under excessive GAs (Table 3), as previously reported in carrot and *Eucalyptus grandis* [WANG & al. 2015; LIU & al. 2018]. A wider root diameter has also been described in the *arm*, *sabre*, *cobra*, *erh-1*, *pom-pom1* and α -tubulin under-expressing mutants, and in plants exposed to 1-butanol, umbelliferone, gamma irradiation or P deficiency [JANKAY & MÜLLER, 1976; SCHNEIDER & al. 1997; BAO & al. 2001; MA & al. 2001; GARDINER & al. 2003; NAGATA & al. 2004; HERMANS & al. 2010; PIETRA, 2014]. Moreover, the reduction of the actin cytoskeleton induces the radial expansion of plant cells, making them shorter and wider, as the interphase MT determine the direction of plant cell elongation [BALUŠKA & al. 2001; BAO & al. 2001]. This means that the capacity of cells to elongate longitudinally depends on the orientation of the cytoskeletal MT [DUGARDEYN & VAN DER STRAETEN, 2008]. Thus, a reduced expression of the α -tubulin gene in *A. thaliana* seedlings results in an abnormal expansion of the root tip (MZ and EZ), with its diameter increasing dramatically at 8 days after germination [BAO & al. 2001]. De-polymerization of MT by oryzalin or 1-butanol also causes the swelling of the root MZ and EZ in *A. thaliana* seedlings, whereas MT stabilization by Taxol expands the root EZ and DZ [BAO & al. 2001; GARDINER & al. 2003].

Therefore, a connection exists between aberrant orientation of MT and reduced cell elongation, as MT regulate the oriented deposition of cellulose microfibrils that determines the direction of cell elongation [BURK & YE, 2002]. More specifically, MT are essential for anisotropic cell expansion because they direct the insertion of cellulose synthase in the CW and guide the orientation of cellulose microfibrils to a perpendicular position with respect to the growth axis, thereby restricting radial cell expansion [JANKAY & MULLER, 1976; LIN & al. 2013]. Interestingly, mutations in P60 katanin protein, essential for anisotropic cell growth, cause an inappropriate feedback regulation of the *DGL1* gen for GAs biosynthesis [KOMORISONO & al. 2005].

With this respect, a link has been proposed between aberrant orientation of MT, radial cell growth and alteration of the root hair patterning [BALUŠKA & al. 2001; BAO & al. 2001]. Thus, a low expression of the α -tubulin gene (TUA6/AS transgenic lines), mutations that inhibit MT polymerization or drugs that brake the actin MT produce aberrant microtubular structures, expand radially the root tip cells, especially at the epidermis and cortex of the MZ and EZ, and induce ectopic root hairs in 5 day-old *A. thaliana* seedlings [BAO & al. 2001; COLLINGS & al. 2006]. On the other hand, the *erh1* and *erh3* mutants, with an altered root hair patterning, exhibit disorganized MT and radially-enlarged layers of root cortex and endodermis, what suggests a connection between radial cell expansion and root hair initiation [SCHNEIDER &

al. 1997; BOUQUIN & al. 2003; MÜLLER & SCHMIDT, 2004; PIETRA & al. 2015]. Moreover, *erh2* is allelic to *pom-1*, a mutant with abnormally-expanded layers of root epidermis and cortex [SCHNEIDER & al. 1997; PIETRA & al. 2015]. In fact, not only cell length, but also cell width differs between Trichoblasts and Atrichoblasts [LÖFKE & al. 2013]. Deficiencies in Fe or P also induce the swelling of root cortical cells, along with ectopic root hairs [PIETRA & al. 2015]. Another clue about the link among MT, cell expansion and cell fate is illustrated by the mutants *cobra* and *sabre*, both with an abnormal cell expansion at the root tip and ectopic root hairs [SCHIEFELBEIN & al. 1997]. The mutation of the SABRE protein, involved in MT organization, causes an abnormal cell expansion at the root cortex [BENFEY & al. 1993], whereas the mutation of the COBRA protein, which is associated to the longitudinal CW of the rapidly-growing root EZ, entails a cellulose deficiency and causes the swelling of the root epidermis and cortex [SCHERES & al. 2002]. Interestingly, the GAs influence CW growth in mesocotyl epidermal cells [PERAZZA & al. 1998]. Therefore, under the experimental conditions of the present study, excessive levels of DELLAs might have impaired the biosynthesis, organization and/or homeostasis of MT in root tip cells of *A. thaliana* seedlings, and this, in turn, might have caused the inhibition of cell elongation and the altered patterning of *GL2* gene expression and of root hairs at the MZ and EZ of the root. In fact, the DELLAs destabilize the MT, giving rise to non-polar cell growth [LOCASCIO & al. 2013].

In this respect, it is known that the levels of ploidy exert an important control over cell size, and that cell size and morphology are, in turn, linked to DNA content [KONDOROSI & al. 2001]. Moreover, in many tissues, cell elongation is associated to the endo-reduplication of the DNA (replication without mitosis that occurs before cell elongation, resulting in a logarithmic accumulation of genome copies in each nucleus) [SANZ & al. 2012]. Thus, the earliest morphological signs in trichome initiation are the induction of endo-reduplication and the increase in the nuclear and cellular size [PERAZZA & al. 1998]. With this regard, it is known that GAs induce endo-reduplication in a dose-dependent manner and regulate cyclin gene expression. In fact, trichomes in the *spy5* mutant have two times more DNA than WT trichomes [PERAZZA & al. 1998; KONDOROSI & al. 2001]. Moreover, in GA-deficient transgenic plants, the observed root swelling, due to MT disorganization, is associated to the induction and accumulation of cyclin CYC3;1 and CYCB1;1 proteins, because the DELLAs are involved in cell cycle progression [UBEDA-TOMÁS & al. 2009; GOU & al. 2010; SÁNCHEZ-CALDERÓN & al. 2013]. In addition, the halting degree of the cell cycle is related to the GA endogenous level [LI & al. 2015b]. Mutations in the α -tubulin gene and drugs that inhibit MT polymerization also induce multi-nucleated cells [BAO & al. 2001]. Interestingly, the control of endo-reduplication in trichomes participates in the regulation of epidermal patterning [PIETRA & al. 2015], whereas the *RHL* genes, related to endo-reduplication, affect root epidermal cell fate independently from the *GL2* gene network [GUO & al. 2009].

Given that stress reduces root cell length and, thus, root length [DINNENY & al. 2008], then, the morphological alterations that were observed in the root cells under excessive levels of DELLAs in 5-day-old *A. thaliana* seedlings might be in tune with the known role of these proteins as mediators of the plant Stress-Induced Morphogenic Responses (SIMR), which are characterized by changes in MT metabolism, CW flexibility and cell cycle progression [POTTERS & al. 2007]. Moreover, it is known that stress inhibits growth by reducing GA levels and promoting the stabilization and accumulation of DELLAs [ACHARD & GENSHIK, 2009; ALONSO-RAMÍREZ & al. 2009], and that the DELLAs mediate the SIMR associated to P deficiency [JIANG & al. 2007]. From this, it might be hypothesized that an alteration of MT homeostasis might have been implicated in the cell size changes that were observed in roots of

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

A. thaliana seedlings grown under (or harbouring) excessive levels of GAs/DELLAs. While excessive levels of DELLAs might have disorganized the MT cytoskeleton, excessive levels of GAs might have stabilized it, giving rise to the changes of cell size and cell fate observed at the MZ and EZ of the root.

Results also suggest that cell fate decisions at the root epidermis might be synchronized with the cell size changes at the inner tissues of the root, such as the cortex. Thus, another reason for the appearance of extra root hairs in *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs/DELLAs might have been the reduction of the ratios of cortical/epidermal cell length (e.g. under excessive DELLAs, which decrease the cortex cell length) and of cortical/epidermal cell width (e.g. under excessive GAs, which decrease the cortex cell width), as they might give rise to epidermal cells laying over two cortical intersections instead of one, and, hence, to two-haired epidermal cells. This would imply that not only the length and width of epidermal cells, but also the length and width of cortical cells might contribute to the number of hairs produced by the root, although additional studies are needed to confirm this hypothesis.

On the other hand, results of this study also point at the cortex, endodermis and pericycle as root tissues from which the GAs/DELLAs might influence the root hair patterning, because transgenic lines over-expressing *gai-1* at these root tissues produced ectopic root hairs and non-hairs [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, it has been shown that blocking GAs signalling at the root endodermis induces morphological defects in the root epidermal cells [LÖFKE & al. 2013; PIETRA & al. 2015; JANES & al. 2018].

Root cell number: Connections to the root hair patterning and abundance

Excessive levels of GAs/DELLAs also altered the radial cell organization in root tips of seedlings of *A. thaliana*. While excessive levels of DELLAs frequently induced additional cells at the epidermis, cortex, endodermis and pericycle of the root, excessive levels of GAs sometimes induced extra cells at the root cortex and pericycle. With this respect, whether the cell proliferation at the root cortex-endodermis-pericycle under excessive levels of DELLAs was another reason for the observed disorganization in the root hair patterning [MCCARTHY-SUÁREZ, unpublished manuscript], it might be worth confirming in future experiments by using inhibitors and/or mutants of cell division.

Regarding the epidermis, interestingly, the predicted number of cells at the Atrichoblast position per root radial section increased under excessive DELLAs (PAC, *gai-1*), but decreased under excessive GAs (*5X* mutant) [MCCARTHY-SUÁREZ, unpublished manuscript]. Nevertheless, as the predicted number of cells at the Trichoblast position did not change, and the percentage of ectopic root hairs was higher under excessive DELLAs as compared to excessive GAs [MCCARTHY-SUÁREZ, unpublished manuscript], then, the higher abundance of root hairs seen under excessive DELLAs in comparison to excessive GAs was probably due to the induction of ectopic root hairs – and, thus, to the higher number of cells at the Atrichoblast position – and not because of the appearance of new Trichoblast positions, given that the number of cells at the Trichoblast position remained unchanged [MCCARTHY-SUÁREZ, unpublished manuscript].

In fact, *Arabidopsis* increases root hair density in the radial dimension by increasing the number of epidermal cells that differentiate into root hair cells [JANES & al. 2018]. An increased number of epidermal cells in the radial domain of the root has also been described under P deficiency and in *tip1* mutants [MA & al. 2001; GRIERSON & SCHIEFELBEIN, 2002; MÜLLER & SCHMIDT, 2004]. However, under P deficiency, the extra epidermal cells at the

Trichoblast position (up to 12) do not increase the abundance of root hairs in the radial axis as ectopic non-hairs also appear [MA & al. 2001; JANES & al. 2018]. Interestingly, a distorted radial patterning of root cells has also been described in mutants of WRKY75, a negative regulator of root hair formation [RISHMAWI & al. 2014].

Radial proliferation of the root cortex cells has also been reported under stress (e.g. P deficiency) as well as in *tip1*, *erh3* and *jkd* mutants, all with an altered root hair patterning [MA & al. 2001; MÜLLER & SCHMIDT, 2004; HASSAN & al. 2010; CUI, 2015; JANES & al. 2018]. Periclinal cell divisions (extra layers) of the root cortex have equally been described in mutants of JKD, which acts from the root cortex to specify the patterning of epidermal cell types [WELCH & al. 2007; LYER-PASCUZZI & BENFEY, 2008; HASSAN & al. 2010]. Interestingly, the GAs restrict the production of extra layers of cortex cells in *Medicago truncatula* roots, thereby generating thinner roots [FONOUNI-FARDE & al. 2019]. In contrast, PAC treatments, or mutations in components of GA signalling, increase the number of layers of root cortex cells, that is, they induce a premature middle cortex (MC) [PAQUETTE & BENFEY, 2005; CUI & BENFEY, 2009]. Moreover, the GAs suppress the MC formation that is proper of the root responses to stress, whereas the DELLAs promote it [CUI & BENFEY, 2009; FONOUNI-FARDE & al. 2019]. Thus, the formation of a MC, due to random and periclinal cell divisions at the root endodermis, and that later on will acquire identity of root cortex, has been described under P deficiency [CUI & BENFEY, 2009; JANES & al. 2018]. Although the production of a MC has also been reported in roots of 3-day-old WT *A. thaliana* seedlings, the presence of a premature MC in the *spy* mutant, with high levels of GAs, suggests that imbalances in GAs/DELLAs homeostasis, which can be triggered by stress, might bring about the formation of a MC [CUI & BENFEY, 2009; CUI, 2015]. Maybe, this was the reason, in this study, for the presence of a MC in roots of *A. thaliana* seedlings grown under (or harbouring) excessive levels of GAs (1 μ M GA₄ and *QD*) (Figures 5 and 6).

Because the PAC-inducible phenotype of MC is also present in the *scr* (scarecrow) and HDA mutants, as well as in trichostatin A (TSA)-treated plants, all producing ectopic root hairs [CUI & BENFEY, 2009], then, a possible link between MC formation (or ectopic cell proliferation at the cortex/endodermis) and alteration of the root hair patterning might be established. Whether this contributed to the disorganisation of the root hair patterning observed under excessive levels of DELLAs, where a cell proliferation was equally observed at the cortex/endodermis/pericycle of the root, it is not known, but might be worth studying in future experiments by using cell division inhibitors and/or mutants. In fact, the alteration of the root hair patterning in the *SCR::gai-1:GR* mutant after growth in DEXA [MCCARTHY-SUÁREZ, unpublished manuscript] was accompanied by random and periclinal cell divisions at the root endodermis (Figure 3). Moreover, because the root MZ and EZ constitute cell fate-decision zones in *Arabidopsis*, then, any changes in cell division at tissues placed underneath the epidermis of the MZ/EZ might bring about changes in epidermal cell fate. In fact, the proliferation of cortex cells is known to influence the root epidermal patterning [LÖFKE et al. 2013; PIETRA & al. 2015; JANES & al. 2018]. Interestingly, histone deacetylation, which affects the root hair patterning, has a role in the proliferation of root cortex cells [XU & al. 2005; LI & al. 2015a]. Increases in the number of root endodermal cells have also been reported in *erh* mutants, during P deficiency, and in *rhizobium*-infected plants [MÜLLER & SCHMIDT, 2004; MA & al. 2001; JANES & al. 2018]. The *schizoriza* (*scz*) mutant, in turn, has defects in the root radial patterning, with extra periclinal cell divisions that result in multiple layers of ground tissue (cortex and endodermis) [MYLONA & al. 2002]. Thus, the results of this study suggest that the alterations in the root hair patterning of *A. thaliana* seedlings grown under

excessive levels of GAs/DELLAs might also have been related to changes in the number of the cortical/endodermal/pericycle cells of the root.

Other possible cause, in this study, for the appearance of ectopic root hairs under excessive levels of GAs/DELLAs [MCCARTHY-SUÁREZ, unpublished manuscript] might have been the anticlinal, diagonal or asymmetric cell divisions (T-clones) frequently observed at the root epidermis, as they gave rise to changes in the *GL2pro::GUS* expression patterning and the size of daughter cells (Figure 3). These T-clones, in turn, might have been linked to alterations in the MT cytoskeleton, as MT are required for the correct positioning of cell division planes [SCHERES & BENFEY, 1999; BAO & al. 2001; RODRIGUEZ-SERRANO & al. 2014]. Moreover, a reduced expression of the α -tubulin gene impairs cell division and results in defects of tissue organization at the root tip [BAO & al. 2001]. Also, the regulation of asymmetric cell divisions in plants is necessary for the generation of cell diversity and patterns [PERNAS & al. 2010]. In fact, root hair cells are shorter than root non-hair cells, so that when an asymmetric cell division takes place at the epidermis of the MZ, the larger cell becomes the root non-hair cell [SALAZAR-HENAO & al. 2016]. With this respect, it is known that the GAs induce cell proliferation at the root MZ and promote the division of epidermal cells [UBEDA-TOMÁS & al. 2009; LEE & al. 2012]. Furthermore, the DELLAs inhibit root cell division in the longitudinal dimension when mediating the SIMR associated to P deficiency [JIANG & al. 2007; PÉRET & al. 2011]. Interestingly, in a dwarf GA-deficient mutant, the MT exhibit an oblique orientation [BOUQUIN & al. 2003]. Also, in the *erh3* mutants, which act in the same route as *cpc* and *rhd6* (root hair defective 6), but independently from WER, the CW are disaligned, diagonally orientated, and malformed, that is, the positioning of the cell plates and the CW is abnormal, what indicates that ERH3 participates in orienting the cell plates during cytokinesis [WEBB & al. 2002]. The cortex-associated SABRE protein is also involved in the orientation of cell division planes [PIETRA, 2014; PIETRA & al. 2013, 2015]. In addition, GL2 is involved in the production of T-clones, whereas WER regulates cell proliferation, what suggests that genes that regulate the cell specification also regulate the cell division planes [LEE & SCHIEFELBEIN, 1999; SCHERES & al. 2002].

Thus, these results suggest that changes in cell number induced by excessive levels of GAs/DELLAs at the epidermis, cortex, endodermis and pericycle of the root tip in *A. thaliana* seedlings might influence the root hair patterning. Moreover, changes in cell number at tissues placed underneath the root epidermis might bring changes of cell fate at the root epidermal cells.

In addition, results of this study suggest that excessive levels of DELLAs in roots of *A. thaliana* seedlings might impair the biosynthesis and/or the assembly of MT, as judged by the swelling of the root tip cells (Figures 1-4), the presence of multi-nucleated cells at the MZ (Figure 6), and the occurrence of ectopic root hairs, branched root hairs, and cells with multiple root hairs [MCCARTHY-SUÁREZ, unpublished manuscript]. In fact, a reduced expression of the α -tubulin gene results in the disassembly and aberrant reorganization of MT [BAO & al. 2001]. This means that the alteration of the root hair patterning in *A. thaliana* seedlings by excessive levels of DELLAs might have been correlated to their inhibitory effect on MT organisation. In fact, MT are essential to establish root cell identity in *Arabidopsis* [WEBB & al. 2002].

The GAs/DELLAs might regulate the root architecture in *A. thaliana* seedlings

Excessive levels of DELLAs also promoted the outburst of LR near the root tip in *A. thaliana* seedlings, whereas excessive levels of GAs inhibited it. This means that any alteration in the levels of GAs/DELLAs might affect not only the root hair patterning, morphology and abundance, but also the root architecture. Thus, in seedlings of *A. thaliana*, physiologically-controlled levels of GAs/DELLAs might have a function in establishing a correct patterning and morphology of root hairs as well as in organising a proper root structure. In fact, supra-physiological levels of DELLAs mediate the root architecture changes that are associated to abiotic stress in plants (i.e., root elongation inhibition, root radial expansion, MC formation, pericycle cell proliferation and LR induction) [YIH & CLARK, 1965; JIANG & al. 2007; GOU & al. 2010; MARTÍN-REJANO & al. 2011; PÉRET & al. 2011; CUI, 2015; WILD & al. 2016]. For example, soil deficiencies of P, B, Fe or NO₃⁻ stimulate LR production, as nutrient concentration regulates LR production [YIH & CLARK, 1965; HERMANS & al. 2010; ZHANG & al. 2014]. Early production of LR has also been reported in the *erh1*, *jkd* and *arm* mutants, equally with a shortened primary root [SCHNEIDER & al. 1997; WELCH & al. 2007]. Interestingly, LR formation, which initiates at the pericycle, is correlated to an alteration in actin and tubulin expression [PASTERNAK & al. 2005; PÉRET & al. 2011; SÁNCHEZ-CALDERÓN & al. 2013]. Therefore, promoting LR outburst by increasing the local levels of DELLAs in roots might constitute a mechanism used by plants to increase the specific area of the root per mass unit, in a similar way as branched root hairs do.

Conclusions

As previously reported for other hormones, the results of this study, and of a previous one [MCCARTHY-SUÁREZ, unpublished manuscript] point to a possible role for the GAs/DELLAs in mediating the changes in the distribution, shape and frequency of root hairs, as well as in the configuration of the root, that take place in plants under situations of stress. Moreover, it is known that auxins, ET, ABA, BRs and SLs mediate these changes without altering the quantitative expression of *WER* and *GL2* [SCHIEFELBEIN, 2003; YANG & al. 2007; MARTÍN-REJANO & al. 2011]. This implies that, by regulating the elongation and/or division of root cells, as well as the production of LR, the GAs/DELLAs might contribute to mediating the changes in the patterning, morphology and abundance of root hairs, and the root architecture adaptations, that occur in plants under environmental stress conditions.

Notes on contributors

Iva MCCARTHY-SUÁREZ – is a postdoctoral researcher in plant biology with special interest in the mechanism of action of plant hormones, senescence and environmental stress.

Acknowledgements

Special thanks to Dr. Miguel Ángel Blázquez and Dr. David Alabadí for supporting the writing of this paper. This study was performed at the blázquez-alabadí Laboratory (Hormone Signalling and Plant Plasticity Group, Instituto de Biología Celular y Molecular de Plantas (IBMCP)-CSIC-UPV, Valencia, Spain). Thanks also to Mrs. Marisol Gascón Irún for confocal microscope technical support. I. McCarthy Suárez also acknowledges a JAE-Doc postdoctoral fellowship (2008-2011), CSIC, Valencia, Spain. This paper is dedicated to the loving memory of Dr. Francisco Culiáñez-Macià, Principal Investigator at the IBMCP-UPV-CSIC, Valencia, Spain.

References

- ACHARD P. & GENSHIK P. 2009. Releasing the brakes of plant growth: how GAs shutdown DELLA proteins. *Journal of Experimental Botany*. **60**(4): 1085-1092. <https://doi.org/10.1093/jxb/ern301>
- ALONSO-RAMÍREZ A., RODRÍGUEZ D., REYES D., JIMÉNEZ J. A., NICOLÁS G., LÓPEZ-CLIMENT M., GÓMEZ-CÁRDENAS A. & NICOLÁS C. 2009. Evidence for a role of gibberellins in salicylic acid-modulated early plant responses to abiotic stress in *Arabidopsis* seeds. *Plant Physiology*. **150**(3): 1335-1344. <https://doi.org/10.1104/pp.109.139352>
- BAHIN E., BAILLY C., SOTTA B., KRANNER I., CORBINEAU F. & LEYMARIE J. 2011. Crosstalk between reactive oxygen species and hormonal signalling pathways regulates grain dormancy in barley. *Plant, Cell & Environment*. **34**(6): 980-993. <https://doi.org/10.1111/j.1365-3040.2011.02298.x>
- BALUŠKA F., JASIK J., EDELMANN H. G., SALAJOVÁ T. & VOLKMANN D. 2001. Latrunculin B-induced plant dwarfism: plant cell elongation is F-actin dependent. *Developmental Biology*. **231**: 113-124. <https://doi.org/10.1006/dbio.2000.0115>
- BAND L. R., UBEDA-TOMÁS S., DYSON R. J., MIDDLETON A. M., HODGMAN C., OWEN M. R., JENSEN O. E., BENNETT M. J. & KING J. R. 2012. Growth-induced hormone dilution can explain the dynamics of plant root cell elongation. *Proceedings of the National Academy of Sciences*. **109**(19): 7577-7582. <https://doi.org/10.1073/pnas.1113632109>
- BAO Y., KOST B. & CHUA N. H. 2001. Reduced expression of α -tubulin genes in *Arabidopsis thaliana* specifically affects root growth and morphology, root hair development and root gravitropism. *Plant Journal*. **28**(2): 145-157. <https://doi.org/10.1046/j.1365-313x.2001.01142.x>
- BENFEY P. N., LINSTED P. J., ROBERTS K., SCHIEFELBEIN J. W., HAUSER M. T. & AESBACHER R. A. 1993. Root development in *Arabidopsis*: Four mutants with dramatically altered root morphogenesis. *Development*. **119**(1): 57-70.
- BOUQUIN T., MATTSON O., NAESTED H., FOSTER R. & MUNDY J. 2003. The *Arabidopsis lue1* mutant defines a katanin p60 ortholog involved in hormonal control of microtubule orientation during cell growth. *Journal of Cell Science*. **116**(Pt 5): 791-801. <https://doi.org/10.1242/jcs.00274>
- BURK D. H. & YE Z. H. 2002. Alteration of oriented deposition of cellulose microfibrils by mutation of a katanin-like microtubule severing protein. *Plant Cell*. **14**(9): 2145-2160. <https://doi.org/10.1105/tpc.003947>
- CAO X. F., LINSTED P., BERGER F., KIEBER J. & DOLAN L. 1999. Differential ethylene sensitivity of epidermal cells is involved in the establishment of cell pattern in the *Arabidopsis* root. *Physiologiae Plantarum*. **106**(6): 311-317. <https://doi.org/10.1034/j.1399-3054.1999.106308.x>
- CHEN C. Y., WU K. & SCHMIDT W. 2015. The histone deacetylase HDA19 controls root cell elongation and modulates a subset of phosphate starvation responses in *Arabidopsis*. *Science Reports*. **5**: 15708. <https://doi.org/10.1038/srep15708>
- CHIEN J. C. & SUSSEX I. M. 1996. Differential regulation of trichome formation on the adaxial and abaxial surfaces by gibberellins and photoperiod in *Arabidopsis thaliana* (L.) Heynh. *Plant Physiology*. **111**(4): 1321-1328. <https://doi.org/10.1104/pp.111.4.1321>
- COLLINGS D. A., LILL A. W., HIMMELSPACH R. & WASTENEYS G. O. 2006. Hypersensitivity to cytoskeletal antagonists demonstrates microtubule-microfilament cross-talk in the control of root elongation in *Arabidopsis thaliana*. *New Phytologist*. **170**(2): 275-290. <https://doi.org/10.1111/j.1469.8137.2006.01671.x>
- CUI H. 2015. Cortex proliferation in the root is a protective mechanism against abiotic stress. *Plant Signalling & Behaviour*. **10**(5): e1011949. <https://doi.org/10.1080/15592324>
- CUI H. & BENFEY P. 2009. Interplay between scarecrow, GA and like heterochromatin protein 1 in ground tissue patterning in the *Arabidopsis* root. *Plant Journal*. **58**(6): 1016-1027. <https://doi.org/10.1111/j.1365-313X.2009.03839.x>
- DINNENY J. R., LONG T. A., WANG J. Y., JUNG J. W., MACE D., POINTER S., BARRON C., BRADY S. M., SCHIEFELBEIN J. & BENFEY P. N. 2008. Cell identity mediates the responses of *Arabidopsis* roots to abiotic stress. *Science*. **320**(5878): 942-945. <https://doi.org/10.1126/science.1153795>
- DUGARDEYN J. & VAN DER STRAETEN D. 2008. Ethylene: Fine tuning plant growth and development by stimulation and inhibition of elongation. *Plant Science*. **175**(1-2): 59-70. <https://doi.org/10.1016/j.plantsci.2008.02.003>
- FONOUNI-FARDE C., MIASSOD A., LAFFONT C., MORIN H., BENDAHMANE A., DIET A. & FRUGIER F. 2019. Gibberellins negatively regulate the development of *Medicago truncatula* root system. *Science Reports*. **9**: 2335. <https://doi.org/10.1038/s41598-019-38876-1>
- GARDINER J., COLLINGS D. A., HARPER J. D. I. & MARC J. 2003. The effects of the phospholipase D- antagonist 1-butanol on seedlings development and microtubule organisation in *Arabidopsis*. *Plant Cell Physiology*. **44**(7): 687-696. <https://doi.org/10.1093/pcp/pcg095>

- GENDRE D., BARAL A., DANG X., ESNAY N., BOUTTÉ Y., STANISLAS T., VAIN T., CLAVEROL S., GUSTAVSSON A., LIN D., GREBE M. & BHALERAO R. P. 2019. Rho-of-plant activated root hair formation requires *Arabidopsis YIP4a/b* gene function. *Development*. **146**(5): dev168559. <https://doi.org/10.1242/dev.168559>
- GOU J., STRAUSS S. H., TSAI C. J., FANG K., CHEN Y., JIANG X. & BUSOV V. B. 2010. Gibberellins regulate lateral root formation in *Populus* through interactions with auxin and other hormones. *Plant Cell*. **22**(3): 623-639. <https://doi.org/10.1105/tpc.109.073239>
- GRIERSON C. & SCHIEFELBEIN J. 2002. Root hairs. pp 2-22. In: SOMERVILLE C. R. & MEYEROWITZ E. M. (eds). *The Arabidopsis book*. American Society of Plant Biologists, Rockville, MD.
- GUO K., KONG W. W. & YANG Z. M. 2009. Carbon monoxide promotes root hair development in tomato. *Plant Cell & Environment*. **32**(8): 1033-1045. <https://doi.org/10.1111/j.1365-3040.2009.01986.x>
- HASSAN H., SCHERES B. & BLILOU I. 2010. Jackdaw controls epidermal patterning in the *Arabidopsis* root meristem through a non-cell autonomous mechanism. *Development*. **137**(9): 1523-1529. <https://doi.org/10.1242/dev.048777>
- HERMANS C., PORCO S., VERBRUGGEN N. & BUSH D. R. 2010. Chitinase-like protein CTL-1 plays a role in altering root system architecture in response to multiple environmental conditions. *Plant Physiology*. **152**(2): 904-917. <https://doi.org/10.1104/pp.109.149849>
- JANES G., VON WANGENHEIM D., COWLING S., KERR I., BAND L., FRENCH A. P. & BISHOP A. 2018. Cellular patterning of *Arabidopsis* roots under low phosphate conditions. *Frontiers in Plant Science*. **9**: 735. <https://doi.org/10.3389/fpls.2018.00735>
- JANKAY P. & MULLER W. H. 1976. The relationship among umbelliferone, growth, and peroxidase levels in cucumber roots. *American Journal of Botany*. **63**(1): 126-132. <https://doi.org/10.2307/2441675>
- JIANG C., GAO X., LIAO L., HARBERD N. P. & FU X. 2007. Phosphate starvation, root architecture and anthocyanin accumulation responses are modulated by the gibberellin-DELLA signalling pathway in *Arabidopsis*. *Plant Physiology*. **145**(4): 1460-1470. <https://doi.org/10.1104/pp.107.103788>
- KAPPUSAMY K. T., CHEN A. Y. & NEMHAUSER J. L. 2009. Steroids are required for epidermal cell fate establishment in *Arabidopsis* roots. *Proceedings of the National Academy of Sciences*. **106**(19): 8073-8076. <https://doi.org/10.1073/pnas.0811633106>
- KOMORISONO M., UEGUCHI-TANAKA M., AICHI I., HASEGAWA Y., ASHIKARI M., KITANO H., MATSUOKA M. & SAZUKA T. 2005. Analysis of the rice mutant *dwarf* and *gladius leaf1*. Aberrant katanin-mediated microtubule organization causes up-regulation of gibberellin biosynthetic genes independently of gibberellin signalling. *Plant Physiology*. **138**(4): 1982-1993. <https://doi.org/10.1104/pp.105.062968>
- KONDOROSI E., ROUDIER F. & GENDREAU E. 2001. Plant cell size control: growing by ploidy? *Current Opinion in Plant Biology*. **3**(6): 488-492. [https://doi.org/10.1016/s1369-5266\(00\)00118-7](https://doi.org/10.1016/s1369-5266(00)00118-7)
- LEE M. M. & SCHIEFELBEIN J. 1999. WEREWOLF, a MYB-related protein in *Arabidopsis*, is a position-dependent regulator of epidermal cell patterning. *Cell*. **99**(5): 473-483. [https://doi.org/10.1016/s0092-8674\(00\)81536-6](https://doi.org/10.1016/s0092-8674(00)81536-6)
- LEE L. Y., HOU X., FANG L., FANG S., KUMAR P. P. & YU H. 2012. Stunted mediates the control of cell proliferation by GA in *Arabidopsis*. *Development*. **139**(9): 1568-1576. <https://doi.org/10.1242/dev.079426>
- LI D. X., CHEN W. Q., XU Z. H. & BAI S. N. 2015a. HISTONE DEACETYLASE6-Defective mutants show increased expression and acetylation of enhancer of tryptrychon and caprice1 and glabra2 with small but significant effects on root epidermis cellular pattern. *Plant Physiology*. **168**(4): 1448-1458. <https://doi.org/10.1104/pp.15.00821>
- LI J., ZHAO Y., CHU H., WANG L., FU Y., LIU P., UPADHYAYA N., CHEN C., MOU T., FENG Y., KUMAR P. & XU J. 2015b. Shoebox modulates root meristem size in rice through dose-dependent effects of gibberellins on cell organisation and proliferation. *PLoS Genetics*. **11**(8): e1005464. <https://doi.org/10.1371/journal.pgen.1005464>
- LIN D., CAO L., ZHOU Z., ZHU L., EHRHARDT D., YANG Z. & FU Y. 2013. Rho GTPase signalling activates microtubule severing to promote microtubule ordering in *Arabidopsis*. *Current Biology*. **23**(4): 290-297. <https://doi.org/10.1016/j.cub.2013.01.022>
- LIU Q. Y., GUO G. S., QIU Z. F., LI X. D., ZENG B. S. & FAN C. J. 2018. Exogenous GA₃ application altered morphology, anatomic and transcriptional regulatory networks of hormones in *Eucalyptus grandis*. *Protoplasma*. **255**(4): 1107-1119. <https://doi.org/10.1007/s00709-018-1218-0>
- LOCASCIO A., BLÁZQUEZ M. A. & ALABADÍ D. 2013. Dynamic regulation of cortical microtubule organization through prefoldin-DELLA interaction. *Current Biology*. **23**(9): 804-809. <https://doi.org/10.1016/j.cub.2013.03.053>
- LÖFKE C., DÜNSTER K. & KLEINE-VEHN J. 2013. Epidermal patterning genes impose non-cell autonomous cell size determination and have additional roles in root meristem size control. *Journal of Integrative Plant Biology*. **55**(9): 864-875. <https://doi.org/10.1111/jipb.12097>

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

- LOMBARDO M. C., GRAZIANO M., POLACCO J. C. & LAMATTINA L. 2006. Nitric oxide functions as a positive regulator of root hair development. *Plant, Signalling & Behaviour*. **1**(1): 28-33. <https://doi.org/10.4161/psb.1.1.2398>
- LYER-PASCUZZI A. & BENFEY P. N. 2008. Transcriptional networks in root cell fate specification. *Biochimica Biophysica Acta*. **1789**(4): 315-325. <https://doi.org/10.1016/j.bbaggm.2008.09.006>
- MA Z., BIELEBERG G. D., BROWN K. M. & LYNCH J. P. 2001. Regulation of root hair density of phosphorus availability in *Arabidopsis thaliana*. *Plant, Cell & Environment*. **24**(4): 459-467. <https://doi.org/10.1046/j.1365-3040.2001.00695.x>
- MARTÍN-REJANO E. M., CAMACHO-CRISTÓBAL J. J., HERRERA-RODRÍGUEZ M. B., REXACH J., NAVARRO-GOCHICOA M. T. & GONZÁLEZ-FONTES A. 2011. Auxin and ethylene are involved in the responses of root system architecture to low boron supply in *Arabidopsis* seedlings. *Physiologiae Plantarum*. **142**(2): 170-178. <https://doi.org/10.1111/j.1399-3054.2011.01459.x>
- MYLONA P., LINSTEAD P., MARTIENSSEN R. & DOLAN L. 2002. Schizoriza controls an asymmetric cell division and restricts epidermal identity in the *Arabidopsis* root. *Development*. **129**(18): 4327-4334. <https://doi.org/10.1242/dev.129.18.4327>
- MÜLLER M. & SCHMIDT W. 2004. Environmentally induced plasticity of root hair development in *Arabidopsis*. *Plant Physiology*. **134**(1): 409-419. <https://doi.org/10.1104/pp.103.029066>
- NAGATA T., TODORIKI S. & KIKUCHI S. 2004. Radial expansion of root cells and elongation of root hairs of *Arabidopsis thaliana* induced by massive doses of gamma irradiation. *Plant Cell Physiology*. **45**(11): 1557-1565. <https://doi.org/10.1093/pcp/pch178>
- NELISSEN H., RYMEN B., JIKUMARU Y., DEMUYNCK K., VAN LIJSEBETTENS M., KAMIYA Y., INZÉ D. & BEEMSTER G. T. S. 2012. A local maximum in gibberellin levels regulates maize leaf growth by spatial control of cell division. *Current Biology*. **22**(13): 1183-1187. <https://doi.org/10.1016/j.cub.2012.04.065>
- NIU Y., JIN C., JIN G., ZHOU Q., LIN X., TANG C. & ZHANG Y. 2011. Auxin modulates the enhanced development of root hairs in *Arabidopsis thaliana* (L.) Heyhn. under elevated CO₂. *Plant, Cell & Environment*. **34**(8): 1304-1317. <https://doi.org/10.1111/j.1365-3040.2011.02330.x>
- OHASHI Y., OKA A., RODRÍGUEZ-POUSADA R., POSSENTI M., RUBERTI I., MORELLI G. & AOYAMA T. 2003. Modulation of phospholipid signalling by GLABRA2 in root hair pattern formation. *Science*. **300**(5624): 1427-1430. <https://doi.org/10.1126/science.1083695>
- PAQUETTE A. J. & BENFEY P. N. 2005. Maturation of the ground tissue of the root is regulated by gibberellin and SCARECROW and requires SHORT-ROOT. *Plant Physiology*. **138**(2): 636-640. <https://doi.org/10.1104/pp.104.058362>
- PASTERNAK T., POTTERS G., CAUBERGS R. & JANSEN M. A. K. 2005 Complementary interactions between oxidative stress and auxins control plant growth responses at plant, organ and cellular level. *Journal of Experimental Botany*. **56**(418): 1991-2001. <https://doi.org/10.1093/jxb/eri196>
- PERAZZA D., VACHON G. & HERZOG M. 1998. Gibberellins promote trichome formation by up-regulating GLABROUS1 in *Arabidopsis*. *Plant Physiology*. **117**(2): 375-383. <https://doi.org/10.1104/pp.117.2.375>
- PÉRET B., CLÉMENT M., NUSSAUME L. & DESNOS T. 2011. Root developmental adaptation to phosphate starvation: Better safe than sorry. *Trends in Plant Sciences*. **16**(8): 442-450. <https://doi.org/10.1016/j.tplants.2011.05.006>
- PERNAS M., RYAN E. & DOLAN L. 2010. Schizoriza controls tissue system complexity in plants. *Current Biology*. **2**(9): 818-823. <https://doi.org/10.1016/j.cub.2010.02.062>
- PIETRA S. 2014. *Characterization of new players in planar polarity establishment in Arabidopsis*. PhD thesis. Umea Plant Science Centre Fysiologisk Botanik, Sweden.
- PIETRA S., GUSTAVSSON A., KIEFER C., KALMBACH L., HÖRSTEDT P., IKEDA Y., STEPANOVA A. N., ALONSO J. M. & GREBE M. 2013. *Arabidopsis* SABRE and CLASP interact to stabilize cell division plane orientation and planar polarity. *Nature Communications*. **4**: 2779. <https://doi.org/10.1038/ncomms3779>
- PIETRA S., LANG P. & GREBE M. 2015. SABRE is required for stabilization of root hair patterning in *Arabidopsis thaliana*. *Physiologiae Plantarum*. **153**(3): 440-453. <https://doi.org/10.1111/ppl.12257>
- POTTERS G., PASTERNAK T. P., GUISEZ Y., PALME K. J. & JANSEN M. A. K. 2007 Stress-induced morphogenic responses: Growing out of trouble? *Trends in Plant Science*. **12**(3): 98-105. <https://doi.org/10.1016/j.tplants.2007.01.004>
- RISHMAWI L., PESCH M., JUENGST C., SCHAUSS A. C., SCHRADER A. & HÜLSKAMP M. 2014. Non-cell autonomous regulation of root hair patterning genes by WRKY75 in *Arabidopsis*. *Plant Physiology*. **165**(1): 186-195. <https://doi.org/10.1104/pp.113.233775>
- RODRÍGUEZ-SERRANO M., PAZMIÑO D. M., SPARKES I., ROCHETTI A., HAWES C., ROMERO-PUERTAS M. C. & SANDALIO L. M. 2014. 2,4-dichlorophenoxyacetic acid promotes S-nitrosylation and oxidation of actin affecting cytoskeleton and peroxisomal dynamics. *Journal of Experimental Botany*. **65**(17): 4783-4793. <https://doi.org/10.1093/jxb/eru237>

- SALAZAR-HENAO J. E., VÉLEZ-BERMÚDEZ I. C. & SCHMIDT W. 2016. The regulation and plasticity of root hair patterning and morphogenesis. *Development*. **143**(11): 1848-1858. <https://doi.org/10.1242/dev.132845>
- SÁNCHEZ-CALDERÓN L., IBARRA-CORTÉS M. E. & ZEPEDA-JAZO I. 2013. Root development and abiotic stress adaptation. pp. 135-168. In: VAHDATI K. & LESLIE C. (eds). *Abiotic stress-plant responses and applications in agriculture*. Intech Open Science, London.
- SANZ L., MURRAY J. A. H. & DEWITTE W. 2012. To divide and to rule: Regulating cell division in roots during post-embryonic growth. *Progress in Botany*. **73**: 57-80. https://doi.org/10.1007/978-3-642-22746-2_2
- SCHERES B. & BENFEY P. N. 1999. Asymmetric cell division in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*. **50**: 505-537. <https://doi.org/10.1146/annurev.arplant.50.1.505>
- SCHERES B., BENFEY P. & DOLAN L. 2002. Root development. pp. 2-18. In: SOMERVILLE C. R. & MEYEROWITZ E. M. (eds). *The Arabidopsis book*. American Society of Plant Biologists, Rockville, MD.
- SCHIEFELBEIN J. 2003. Cell fate specification in the epidermis: A common patterning mechanism in the root and the shoot. *Current Opinion in Plant Biology*. **6**(1): 74-78. <https://doi.org/10.1016/S136952660200002X>
- SCHIEFELBEIN J., MASUCCI J. D. & WANG H. 1997. Building a root: The control of patterning and morphogenesis during root development. *Plant Cell*. **9**(7): 1089-1098. <https://doi.org/10.1105/tpc.9.7.1089>
- SCHMIDT W., TITTEL J. & SCHIKORA A. 2000. Role of hormones in the induction of iron deficiency responses in *Arabidopsis* roots. *Plant Physiology*. **122**(4): 1109-1118. <https://doi.org/10.1104/pp.122.4.1109>
- SCHNEIDER K., WELLS B., DOLAN L. & ROBERTS K. 1997. Structural and genetic analysis of epidermal cell differentiation in *Arabidopsis* primary roots. *Development*. **124**(9): 1789-1798.
- SHANI E., WEINSTAIN R., ZHANG Y., CASTILLEJO C., KAISERLI E., CHORY J., TSIEN R. Y. & ESTELLE M. 2013. Gibberellins accumulate in the elongating endodermal cells of *Arabidopsis* root. *Proceedings of the National Academy of Sciences*. **110**(12): 4834-4839. <https://doi.org/10.1073/pnas.1300436110>
- SHIN L. J., HUANG H. E., CHANG H., LIN Y. N., FENG T. Y. & GER M. J. 2011. Ectopic ferredoxin I protein promotes root hair growth through induction of reactive oxygen species in *Arabidopsis thaliana*. *Journal of Plant Physiology*. **168**(5): 434-440. <https://doi.org/10.1016/j.jplph.2010.08.002>
- TRAW M. B. & BERGELSON J. 2003. Interactive effects of jasmonic acid, salicylic acid, and gibberellin on induction of trichomes in *Arabidopsis*. *Plant Physiology*. **133**(3): 1367-1375. <https://doi.org/10.1104/pp.103.027086>
- UBEDA-TOMÁS S., SWARUP R., COATES J., SWARUP K., LAPLACE L., BEEMSTER G. T. S., HEDDEN P., BHALERAO R. & BENNETT M. J. 2008. Root growth in *Arabidopsis* requires gibberellin/DELLA signalling in the endodermis. *Nature Cell Biology*. **10**(5): 625-628. <https://doi.org/10.1038/ncb1726>
- UBEDA-TOMÁS S., FEDERICI F., CASIMIRO I., BEEMSTER G. T. S., BHALERAO R., SWARUP R., DOERNER P., HASELHOFF J. & BENNETT M. J. 2009. Gibberellin signalling in the endodermis controls *Arabidopsis* root meristem size. *Current Biology*. **19**(14): 1194-1199. <https://doi.org/10.1016/j.cub.2009.06.023>
- VAN HENGEL A. J., BARBER C. & ROBERTS K. 2004. The expression patterns of arabinogalactan-protein *AtAGP30* and *GLABRA2* reveal a role for abscisic acid in the early stages of root epidermal patterning. *Plant Journal*. **39**(1): 70-83. <https://doi.org/10.1111/j.1365-313X.2004.02104.x>
- WANG G. L., QUE F., XU Z. S., WANG F. & XIONG A. S. 2015. Exogenous gibberellin altered morphology, anatomic and transcriptional regulatory networks of hormones in carrot root and shoot. *BMC Plant Biology*. **15**: 290. <https://doi.org/10.1186/s12870-015-0679-y>
- WEBB M., JOUANNIC S., FOREMAN J., LINSTED P. & DOLAN L. 2002. Cell specification in the *Arabidopsis* root epidermis requires the activity of *ECTOPIC ROOT HAIR 3* - a katanin-P60 protein. *Development*. **129**(1): 123-131.
- WELCH D., HASSAN H., BLILOU I., IMMINK R., HEIDSTRA R. & SCHERES B. 2007. *Arabidopsis* JACKDAW and MAGPIE zinc finger proteins delimit asymmetric cell division and stabilize tissue boundaries by restricting SHORT-ROOT action. *Genes Development*. **21**(17): 2196-2204. <https://doi.org/10.1101/gad.440307>
- WILD M., DAVIÈRE J. M., REGNAULT T., SAKVARELIDZE-ACHARD L., CARRERA E., LOPEZ-DIAZ I., CAYREL A., DUBEAUX G., VERT G. & ACHARD P. 2016. Tissue-specific regulation of gibberellin signalling fine-tunes *Arabidopsis* iron-deficiency responses. *Developmental Cell*. **37**(2): 190-200. <https://doi.org/10.1016/j.devcel.2016.03.022>
- XU C. R., LIU C., WANG Y. L., LI L. C., CHEN W. Q., XU Z. H. & BAI S. N. 2005. Histone deacetylation affects expression of cellular patterning genes in the *Arabidopsis* root epidermis. *Proceedings of the National Academy of Sciences*. **102**(40): 14469-14474. <https://doi.org/10.1073/pnas.0503143102>
- YANG T., SAVAGE N. & SCHMIDT W. 2007. Plasticity of root epidermal cell fate in response to nutrient starvation. 18th International Conference on *Arabidopsis* Research. TAIR accession publication [online]: 501721882 [accessed Aug. 23rd, 2021].
- YIH R. Y. & CLARK H. E. 1965. Carbohydrate and protein content to boron-deficient tomato root tips in relation to anatomy and growth. *Plant Physiology*. **40**(2): 312-315. <https://doi.org/10.1104/pp.40.2.312>

SUPRA-PHYSIOLOGICAL LEVELS OF GIBBERELLINS / DELLAS MODIFY ROOT CELL SIZE...

ZHANG C., BOUSQUET A. & HARRIS J. M. 2014. Abscisic acid and lateral root organ defective/numerous infections and polyphenolics modulate root elongation via reactive oxygen species in *Medicago truncatula*. *Plant Physiology*. **166**(2): 644-658. <https://doi.org/10.1104/pp.114.248542>

How to cite this article:

McCARTHY-SUÁREZ I. 2021. Supra-physiological levels of gibberellins / DELLAs modify root cell size / number and root architecture in root tips of *A. thaliana* seedlings. Connections to the root hair patterning and abundance. *J. Plant Develop.* **28**: 69-96. <https://doi.org/10.47743/jpd.2021.28.1.887>
